arXiv:1205.6869v1 [math.CO] 31 May 2012

A NEW UPPER BOUND ON THE ACYCLIC CHROMATIC INDICES OF PLANAR GRAPHS

Weifan Wang[*], Qiaojun Shu
Department of Mathematics, Zhejiang Normal University, Jinhua 321004, China

Yiqiao Wang
Academy of Mathematics and Systems Science, Beijing 100080, China

Abstract

An acyclic edge coloring of a graph G is a proper edge coloring such that no bichromatic cycles are produced. The acyclic chromatic index $a'(G)$ of G is the smallest integer k such that G has an acyclic edge coloring using k colors. It was conjectured that $a'(G) \leq \Delta + 2$ for any simple graph G with maximum degree Δ. In this paper, we prove that if G is a planar graph, then $a'(G) \leq \Delta + 7$. This improves a result by Basavaraju et al. [*Acyclic edge-coloring of planar graphs*, SIAM J. Discrete Math., 25 (2011), pp. 463-478], which says that every planar graph G satisfies $a'(G) \leq \Delta + 12$.

Key words. Acyclic edge coloring, planar graph, girth, maximum degree

AMS subject classification. 05C15

1 Introduction

Only simple graphs are considered in this paper. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. A *proper edge k-coloring* is a mapping $c : E(G) \to \{1, 2, \ldots, k\}$ such that any two adjacent edges receive different colors. The *chromatic index* $\chi'(G)$ of G is the smallest integer k such that G is edge k-colorable. A proper edge k-coloring c of G is called *acyclic* if there are no bichromatic cycles in G, i.e., the union of any two color classes induces a subgraph of G that is a forest. The *acyclic chromatic index* of G, denoted by $a'(G)$, is the smallest integer k such that G is acyclically edge k-colorable.

Let $\Delta(G)$ (Δ for short) denote the maximum degree of a graph G and $g(G)$ denote the girth of G, i.e., the length of a shortest cycle in G. By Vizing's Theorem [23], $\Delta \leq \chi'(G) \leq \Delta + 1$. Thus, it is obvious that $a'(G) \geq \chi'(G) \geq \Delta$. Fiamčik [8], and later Alon, Sudakov and Zaks [1], made independently the following conjecture:

Conjecture 1 *For any graph G, $a'(G) \leq \Delta + 2$.*

---

[*] Research supported partially by NSFC(No.11071223) and ZJNSF(No.Z6090150); Corresponding author. Email: wwf@zjnu.cn.

Using probabilistic method, Alon, McDiarmid and Reed [2] proved that $a'(G) \leq 64\Delta$ for any graph G. Molloy and Reed [14] improved this bound to that $a'(G) \leq 16\Delta$. Alon, Sudakov and Zaks [1] proved that there is a constant c such that $a'(G) \leq \Delta + 2$ for a graph G whenever $g(G) \geq c\Delta \log \Delta$. They also confirmed Conjecture 1 for almost all Δ-regular graphs. Něsetřil and Wormald [15] gave an improvement to this result by showing that $a'(G) \leq \Delta + 1$ for a random Δ-regular graph G. The acyclic edge coloring of some special classes of graphs was also considered, including subcubic graphs [4, 16], graphs with maximum degree 4 [12], outerplanar graphs [11, 13], series-parallel graphs [10, 21], and planar graphs [3, 7, 9, 17, 18, 20, 22].

Suppose that G is a planar graph. Fiedorowicz et al. [6] proved that $a'(G) \leq 2\Delta + 29$ and Hou et al. [10] proved that $a'(G) \leq \max\{2\Delta - 2, \Delta + 22\}$. The currently best known upper bound on the acyclic chromatic index of a planar graph G is that $a'(G) \leq \Delta + 12$ by Basavaraju et al. [5]. In this paper, we will improve this result by replacing 12 by 7.

Before showing the main result, we need to introduce some definitions and notations.

Given a graph G, let $d_G(v)$ (or simply $d(v)$) denote the degree of a vertex v in G. A vertex of degree k (at least k, at most k) is called a *k-vertex* (k^+-vertex, k^--vertex). For $k \geq 1$, let $n_k(v)$ ($n_{k^+}(v)$, $n_{k^-}(v)$) denote the number of k-vertices (k^+-vertices, k^--vertices) adjacent to a vertex v in G. Let $\delta(G)$ denote the minimum degree of G.

A *plane* graph is a particular drawing in the Euclidean plane of a certain planar graph. For a plane graph H, we denote its face set by $F(H)$ and the degree of a face $f \in F(H)$ is denoted by $d_H(f)$ (or simply $d(f)$). Similarly, we can define a *k-face*, a *k^+-face*, and a *k^--face*. Furthermore, for $f \in F(H)$, we use $b(f)$ to denote the boundary walk of f and write $f = [u_1u_2 \ldots u_n]$ if $u_1, u_2, \ldots, u_n$ are the vertices on $b(f)$ enumerated in some order, where u_i may be identical to u_j for some $i, j \in \{1, 2, \ldots, n\}$ when G contains cut vertices.

2 A structural lemma

In this section, we establish a structural lemma, which plays an important role in the proof of the main result in Section 3.

Lemma 1 *Let G be a 2-connected planar graph with $\Delta \geq 5$. Then G contains one of the following configurations* **($\mathbf{A}_1$)-($\mathbf{A}_4$)**, *as shown in Fig. 1:*

($\mathbf{A}_1$) *A path uvw with $d(v) = 2$ and $d(u) \leq 9$.*

($\mathbf{A}_2$) *A vertex u with $n_2(u) \geq 1$ and $n_{8^-}(u) \geq d(u) - 8$. Suppose that $u_1, u_2, \ldots, u_{d(u)-1}$, v are the neighbors of u such that $d(u_1) \geq d(u_2) \geq \cdots \geq d(u_{d(u)-1}) \geq d(v) = 2$. Let w be the neighbor of v different from u. For $1 \leq i \leq d(u) - 1$, if $d(u_i) = 2$, then we use x_i to denote the neighbor of u_i different from u. Then at least one of the following cases holds:*

($\mathbf{A}_{2.1}$) *$n_{8^-}(u) \geq d(u) - 7$;*

($\mathbf{A}_{2.2}$) *$n_{8^-}(u) = d(u) - 8$, and $n_2(u) \geq d(u) - 9$.*

($\mathbf{A}_3$) *A 3-vertex u is adjacent to a vertex v such that one of the following holds:*

($\mathbf{A}_{3.1}$) *$d(v) \le 8$;*

($\mathbf{A}_{3.2}$) *$d(v) = 9$ and $uu_2, vu_2 \in E(G)$;*

($\mathbf{A}_{3.3}$) *$d(v) = 10$, $n_{5^-}(v) \ge 5$ and $uu_i, vu_i \in E(G)$, $i = 1, 2$.*

($\mathbf{A}_4$) *Suppose that a vertex v is adjacent to $u, v_2, \cdots, v_{d(v)}$ such that $d(u) \le d(v_2) \le \cdots \le d(v_{d(v)})$. Then at least one of the following cases holds:*

($A_{4.1}$) *$d(v) = 4$, $4 \le d(u) \le 7$, and $d(u) + d(v_2) \le 17$;*

($A_{4.2}$) *$d(v) = 5$, $4 \le d(u) \le 6$, and $d(u) + d(v_2) + d(v_3) \le 18$, or $d(u) = d(v_2) = 6$, $d(v_i) = 7$, $i = 3, 4, 5$ and $uv_5 \in E(G)$.*



Fig. 1: Configurations (A_1)-(A_4) in Lemma 1.

Remarks. In Fig. 1, vertices marked solid points have no edges of G incident to them other than those shown, whereas the other vertices have edges connected to other vertices of G not in the configuration.

Proof. Assume to the contrary that G contains none of the configurations (A_1)-(A_4). Since G is 2-connected, it follows that $\delta(G) \ge 2$. Let G' be the graph obtained by removing all the 2-vertices of G and H is a component of G'. Then H is a connected planar graph and for any vertex $v \in V(H)$, v is of degree at least 3 in G.

In what follows, we assume that H is embedded in the plane. Let $v \in V(H)$. Then $v \in V(G)$. The degree of v in G is simply denoted by $d(v)$, whereas the degree of v in H is denoted by $d_H(v)$. For $k \ge 1$, let $n'_k(v)$ ($n'_{k^+}(v)$, $n'_{k^-}(v)$) denote the number of k-vertices (k^+-vertices, k^--vertices) adjacent to v in H, while $n_k(v)$ ($n_{k^+}(v)$, $n_{k^-}(v)$) denote the number of k-vertices (k^+-vertices, k^--vertices) adjacent to v in G. The number of k-vertices (k^+-vertices) lying on the boundary of a face $f \in F(H)$ is denoted by $n_k(f)$ ($n_{k^+}(f)$). Similarly, let $m_k(v)$ denote the number of k-faces incident to v in H, and let $\delta(f)$ denote the minimum degree of vertices on $b(f)$.

Since G contains no (A_1), there is no 9^--vertex adjacent to a 2-vertex, i.e., if u is a vertex with $3 \leq d(u) \leq 9$, then $d_H(u) = d(u)$. Similarly, if $d(u) \geq 10$ and $n_2(u) = 0$, then $d_H(u) = d(u)$. Since G contains neither $(A_{2.1})$ nor $(A_{2.2})$, every 10^+-vertex u is adjacent to at most $(d(u)-9)$ 2-vertices, which implies that if $d(u) \geq 10$ and $n_2(u) \geq 1$, then $d_H(u) \geq 9$, $d_H(u) = d(u) - n_2(u)$.

Claim 1 *Let* $u \in V(H)$.

(a) $\delta(H) \geq 3$.

(b) *If* $d_H(u) \leq 8$, *then* $d(u) = d_H(u)$.

(c) $n'_i(u) = n_i(u)$ *and* $n_{i^-}(u) = n_2(u) + n'_{i^-}(u) = d(u) - d_H(u) + n'_{i^-}(u)$ *for any* $3 \leq i \leq 8$. *In particular,* $n'_{8^-}(u) = \sum\limits_{3 \leq i \leq 8} n'_i(u) = \sum\limits_{3 \leq i \leq 8} n_i(u) = n_{8^-}(u) - n_2(u)$.

Proof. It is easy to see that (c) holds trivially. To prove (a), assume that H contains a 2^--vertex u. Then $d(u) \geq 10$, $n_{8^-}(u) \geq n_2(u) = d(u) - d_H(u) \geq d(u) - 2$ and hence $(A_{2.1})$ is contained in G, a contradiction.

To prove (b), assume that $d_H(u) \leq 8$ and $d(u) \neq d_H(u)$. It is easy to see that $d(u) \geq 10$, and $n_2(u) = d(u) - d_H(u) \geq d(u) - d_H(u) \geq d(u) - 8$, and hence $(A_{2.1})$ or $(A_{2.2})$ is contained in G, also a contradiction. □

Claim 2 *If* $d_H(u) = 9$ *and* $n'_{8^-}(u) \geq 1$, *then* $d(u) = 9$. *In other words, if* $n'_{8^-}(u) \geq 1$, *then* $d_H(u) = d(u) \leq 9$, *or* $d_H(u) \geq 10$.

Proof. If $d(u) \neq 9$, then $n_2(u) = d(u) - 9$, $n_{8^-}(u) \geq n_2(u) + n'_{8^-}(u) \geq n_2(u) + 1 = d(u) - d_H(u) + 1 = d(u) - 8$ and hence $(A_{2.2})$ is contained in G. □

Claim 3 *If* $d_H(u) \geq 10$ *and* $d(u) \neq d_H(u)$, *then* $n_3(u) + n_4(u) + n_5(u) \leq d_H(u) - 8$.

Proof. Otherwise, assume that $n_3(u) + n_4(u) + n_5(u) \geq d_H(u) - 7$. Then, $n_{8^-}(u) \geq n_2(u)+n_3(u)+n_4(u)+n_5(u) = d(u)-d_H(u)+n_3(u)+n_4(u)+n_5(u) \geq d(u)-d_H(u)+d_H(u)-7 = d(u) - 7$ and hence $(A_{2.1})$ is contained in G. □

Claim 4 *If* $f = [uvw]$ *with* $d_H(v) = 3$, *then* $d_H(u) \geq 10$, $d_H(w) \geq 10$ *and* $n_{10^+}(f) = 2$.

Proof. Since G contains neither $(A_{3.1})$ nor $(A_{3.2})$, we have $d(u) \geq 10$ and $d(w) \geq 10$. Obviously, $n'_{8^-}(u) \geq n'_3(u) \geq 1$ and $n'_{8^-}(w) \geq n'_3(w) \geq 1$. Then, by Claim 2, $d_H(u) \geq 10$, $d_H(w) \geq 10$ and $n_{10^+}(f) = 2$. □

Claim 5 *If* $3 \leq d(u) \leq 8$, *then* $n'_{10^+}(u) = n_{10^+}(u)$.

Proof. Note that $d_H(w) \leq d(w)$ for any $w \in V(H)$. Let v be a vertex of degree at least 10 in G and $uv \in E(G)$. Since $uv \in E(H)$, $n'_{8^-}(v) \geq n_3(v) \geq 1$ and $d_H(v) \geq 10$ by Claim 2. $\square$

Claim 6 *Let u be a 4-vertex. If $n_{7^-}(u) \geq 1$, then $n'_{10^+}(u) = 3$. Otherwise, $n'_{8^+}(u) = 4$.* $\square$

Proof. Since G contains neither (A_1) nor $(A_{3.1})$, $n_{3^-}(u) = 0$. If $n_{7^-}(u) \geq 1$, then $n_i(u) \geq 1$ for some $4 \leq i \leq 7$. Since G contains no $(A_{4.1})$, we have $n_{11^+}(u) = 3$ and $n'_{10^+}(u) = 3$ by Claim 5. Otherwise, $n_{8^+}(u) = 4$ and $n'_{8^+}(u) = 4$. $\square$

Claim 7 *Let u be a 5-vertex with $n_4(u) \geq 1$. If $n_{5^-}(u) \geq 2$, then $n'_{10^+}(u) = 3$.*

Proof. Since G contains neither (A_1) nor $(A_{3.1})$, $n_{3^-}(u) = 0$. Since G contains no $(A_{4.2})$, we have $n_{10^+}(u) = 3$ and $n'_{10^+}(u) = 3$ by Claim 5. $\square$

To derive a contradiction, we make use of the discharging method. First, by Euler's formula $|V(H)| - |E(H)| + |F(H)| = 2$ and the relation $\sum\limits_{u\in V(H)} d_H(u) = \sum\limits_{f\in F(H)} d(f) = 2|E(H)|$, we can derive the following identity.

$$\sum_{u\in V(H)} (2d_H(u) - 6) + \sum_{f\in F(H)} (d(f) - 6) = -12. \tag{1}$$

Next, we define an initial weight function w by $w(u) = 2d_H(u) - 6$ for $u \in V(H)$ and $w(f) = d(f) - 6$ for $f \in F(H)$. It follows from (1) that the total sum of weights is equal to -12. In what follows, we will define some discharging rules and redistribute weights accordingly. Once the discharging is finished, a new weight function w' is produced. However, the total sum of weights is kept fixed when the discharging is in process. Nevertheless, we can show that $w'(x) \geq 0$ for all $x \in V(H) \cup F(H)$. This leads to the following obvious contradiction

$$0 \leq \sum_{x\in V(H)\cup F(H)} w'(x) = \sum_{x\in V(H)\cup F(H)} w(x) = -12 < 0$$

and hence demonstrates that no such counterexample can exist.

A 3-face f is called *bad* if f is incident to a 3-vertex, i.e., $\delta(f) = 3$.

Let y be a vertex in H and $f = [\cdots xyz \cdots]$ be a face incident to x, y, z with $d_H(x) \geq d_H(z)$. Let $\tau(y \to f)$ denote the amount of weight transferred from y to f according to the following defined discharging rules:

(R1) $d_H(y) = 4$. If $n_{7^-}(y) = 0$, then $\tau(y \to f) = \frac{1}{2}$. If $n_{7^-}(y) = 1$, then

$$\tau(y \to f) = \begin{cases} \frac{4}{5}, & \text{if } d_H(z) \leq 7; \\ \frac{1}{5}, & \text{if } d_H(z) \geq 8. \end{cases}$$

(R2) $d_H(y) = 5$. If $n_4(y) \geq 1$, then $\tau(y \to f) = \frac{4}{5}$. Otherwise, $n_4(y) = 0$, we carry out the following subrules:

- If $4 \leq d(f) \leq 5$ and $d_H(z) \leq 8$, then $\tau(y \to f) = \frac{1}{2}$
- If $d(f) = 3$, or $4 \leq d(f) \leq 5$ and $d_H(z) \geq 9$, then

$$\tau(y \to f) = \begin{cases} \frac{7}{5}, \text{ if } d_H(x) = d_H(z) = 5; \\ \frac{6}{5}, \text{ if } d_H(x) = 6 \text{ and } d_H(z) = 5; \\ \frac{13}{14}, \text{ if } d_H(x) = 7 \text{ and } d_H(z) = 5; \\ 1, \text{ if } d_H(x) = 8 \text{ and } d_H(z) = 5; \\ \frac{11}{12}, \text{ if } d_H(x) \geq 9 \text{ and } d_H(z) = 5; \\ 1, \text{ if } d_H(x) = d_H(z) = 6; \\ \frac{6}{7}, \text{ if } d_H(x) = 7 \text{ and } d_H(z) = 6; \\ \frac{3}{4}, \text{ if } d_H(x) = 8 \text{ and } d_H(z) = 6; \\ \frac{2}{3}, \text{ if } d_H(x) \geq 9 \text{ and } d_H(z) = 6; \\ \frac{5}{7}, \text{ if } d_H(x) = d_H(z) = 7; \\ \frac{9}{14}, \text{ if } d_H(x) \geq 8 \text{ and } d_H(z) = 7; \\ \frac{1}{2}, \text{ if } d_H(z) = 8; \\ \frac{1}{3}, \text{ if } d_H(z) \geq 9. \end{cases}$$

(R3) $d_H(y) = k$ and $6 \leq k \leq 9$. Then

$$\tau(y \to f) = \frac{2k-6}{k} = \begin{cases} \frac{4}{3}, \text{ if } k = 9; \\ \frac{5}{4}, \text{ if } k = 8; \\ \frac{8}{7}, \text{ if } k = 7; \\ 1, \text{ if } k = 6. \end{cases}$$

(R4) $d_H(y) \geq 10$. If $4 \leq d(f) \leq 5$, or $d(f) = 3$ and $d_H(z) \geq 6$, then $\tau(y \to f) = 1$. Otherwise, $f = [xyz]$ with $d_H(z) \leq 5$. Then

$$\tau(y \to f) = \begin{cases} \frac{3}{2}, \text{ if } d_H(z) = 3; \\ \frac{7}{5}, \text{ if } d_H(z) = 4 \text{ and } n_{7^-}(z) = 1; \\ \frac{5}{4}, \text{ if } d_H(z) = 4 \text{ and } n_{7^-}(z) = 0; \\ \frac{11}{10}, \text{ if } d_H(z) = 5,\ n_4(z) \geq 1, \text{ and } d_H(x) \geq 10; \\ \frac{5}{4}, \text{ if } d_H(z) = 5,\ n_4(z) \geq 1, \text{ and } 6 \leq d_H(x) \leq 9; \\ \frac{7}{5}, \text{ if } d_H(x) = d_H(z) = 5 \text{ and } n_4(z) \geq 1 \text{ or } n_4(x) \geq 1; \\ \frac{4}{3}, \text{ if } d_H(z) = 5,\ n_4(z) = 0 \text{ and } n_4(x) = 0 \text{ if } d_H(x) = 5. \end{cases}$$

It remains to inspect that $w'(x) \geq 0$ for all $x \in V(H) \cup F(H)$. Let $f \in F(H)$. We consider several cases as follows:

Case 1. $d(f) = 3$.

Let $f = [uvw]$. Then $w(f) = d(f) - 6 = -3$. We have some subcases, depending on the size of $\delta(f)$.

(1.1) Assume that $\delta(f) = 3$. Then $n_{10^+}(f) = 2$ by Claim 4 and (R4), each 10^+-vertex in $b(f)$ gives $\frac{3}{2}$ to f. Hence, $w'(f) = -3 + 2 \times \frac{3}{2} = 0$.

(1.2) Assume that $\delta(f) = 4$ with $d_H(v) = 4$ and $d_H(u) \geq d_H(w)$. If $n_{7^-}(v) = 1$, then $d_H(u) \geq 10$ by Claim 6, $d_H(w) \geq 6$, or $4 \leq d_H(w) \leq 5$ and $n_4(w) \geq 1$. Then, u sends $\frac{7}{5}$ to f by (R4) and w sends at least $\frac{4}{5}$ to f by (R1), (R2), (R3) or (R4). Hence,

$w'(f) \geq -3+2\times\frac{4}{5}+\frac{7}{5}=0$. If $n_{7^-}(v)=0$, then $d_H(u), d_H(w) \geq 8$. Then, each of u, w sends $\frac{5}{4}$ to f by (R3) and (R4) and v sends $\frac{1}{2}$ to f by (R1). Hence, $w'(f) \geq -3+2\times\frac{5}{4}+\frac{1}{2}=0$.

(1.3) Assume that $\delta(f)=5$ with $d_H(v)=5$ and $d_H(u) \geq d_H(w)$. We further have three possibilities:

- $n_5(f)=3$. Since G contains none of $(A1)$, $(A3.1)$ and $(A4.2)$, $n_4(u)=n_4(v)=n_4(w)=0$ and each vertex in $b(f)$ sends $\frac{7}{5}$ to f by (R2). Thus, $w'(f) \geq -3+3\times\frac{7}{5}=\frac{6}{5}$.
- $n_5(f)=2$ and $d_H(w)=5$.

If $n_4(v)=1$ or $n_4(w)=1$, then $d(u) \geq 10$ since G contains no $(A4.2)$ and $d_H(u) \geq 10$ by Claim 2. Hence, each of v and w sends $\frac{4}{5}$ or $\frac{11}{12}$ to f by (R2), u sends $\frac{7}{5}$ to f by (R4) and therefore $w'(f) \geq -3+2\times\frac{4}{5}+\frac{7}{5}=0$. Otherwise, $n_4(v)=n_4(w)=0$. By (R2), (R3) or (R4), if $d_H(u)=6$, then $w'(f) \geq -3+(2\times\frac{6}{5}+1)=\frac{2}{5}$; if $d_H(u)=7$, then $w'(f) \geq -3+(2\times\frac{13}{14}+\frac{8}{7})=0$; if $d_H(u)=8$, then $w'(f) \geq -3+(2\times 1+\frac{5}{4})=\frac{1}{4}$; if $d_H(u) \geq 9$, then $w'(f) \geq -3+(2\times\frac{11}{12}+\frac{4}{3})=\frac{1}{6}$.

- $n_5(f)=1$ and $d_H(u) \geq d_H(w) \geq 6$.

If $n_4(v)=1$, then v sends $\frac{4}{5}$ to f by (R2). If $d_H(w)=6$, then $d(u) \geq 9$ and $d_H(u) \geq 9$ since G contains no $(A4.2)$. Thus, w sends 1 to f and u sends $\frac{4}{3}$ to f by (R3) or u sends $\frac{5}{4}$ to f by (R4). Hence, $w'(f) \geq -3+\min\{\frac{4}{5}+1+\frac{5}{4}, \frac{4}{5}+1+\frac{4}{3}\}=\frac{1}{20}$. Otherwise, $d_H(u) \geq d_H(w) \geq 7$ and each of w and u sends at least $\frac{11}{10}$ by (R3) or (R4). Hence, $w'(f) \geq -3+\frac{4}{5}+2\times\frac{11}{10}=0$. Now assume that $n_4(v)=0$. By (R2), (R3) or (R4), if $d_H(u)=d_H(w)=6$, then $w'(f) \geq -3+(1+2\times 1)=0$; if $d_H(u)=7$ and $d_H(w)=6$, then $w'(f) \geq -3+(\frac{6}{7}+1+\frac{8}{7})=0$; if $d_H(u)=8$ and $d_H(w)=6$, then $w'(f) \geq -3+(\frac{3}{4}+1+\frac{5}{4})=0$; if $d_H(u) \geq 9$ and $d_H(w)=6$, then $w'(f) \geq -3+(\frac{2}{3}+1+\frac{4}{3})=0$; if $d_H(u)=d_H(w)=7$, then $w'(f) \geq -3+(\frac{5}{7}+2\times\frac{8}{7})=0$; if $d_H(u) \geq 8$ and $d_H(w)=7$, then $w'(f) \geq -3+(\frac{9}{14}+\frac{8}{7}+\frac{5}{4})=\frac{1}{28}$; if $d_H(u) \geq 8$ and $d_H(w)=8$, then $w'(f) \geq -3+(\frac{1}{2}+2\times\frac{5}{4})=0$; if $d_H(u) \geq d_H(w) \geq 9$, then $w'(f) \geq -3+(\frac{1}{3}+2\times\frac{4}{3})=0$.

(1.4) Assume that $\delta(f) \geq 6$. Then each vertex in $b(f)$ sends at least 1 to f by (R3) or (R4). Hence, $w'(f) \geq -3+3\times 1=0$.

Case 2. $d(f)=4$.

Let $f=[uvst]$. Then $w(f)=d(f)-6=-2$. If $n_{6^+}(f) \geq 2$, then by (R3) or (R4), each 6^+-vertex in $b(f)$ gives at least 1 to f. Hence, $w'(f) \geq -2+2\times 1=0$. Now we assume that $n_{6^+}(f) \leq 1$. Note that if $\delta(f)=3$, then $n_{9^+}(f) \geq 2$ and if there is a 4-vertex v in $b(f)$ with $n_{7^-}(v)=0$, then $n_{8^+}(f) \geq 2$. Thus it suffice to suppose that $\delta(f) \geq 5$, or $\delta(f)=4$ and each 4-vertex v in $b(f)$ having $n_{7^-}(v)=1$. Assume that $\delta(f)=4$ with $d_H(v)=4$ and $d_H(u) \leq 5 \leq 10 \leq d_H(s)$. It is easy to see that $n_4(u) \geq 1$. By (R1), (R2) and (R4), v sends $\frac{4}{5}$ to f, u sends $\frac{4}{5}$ to f, and s sends 1 to f. Hence, $w'(f) \geq -2+1+2\times\frac{4}{5}=\frac{3}{5}$. Assume that $\delta(f)=5$ with $d(u)=d(v)=d(s)=5$ and $d_H(t) \geq 5$. By (R2), (R3) or (R4), each of u, v, s, t gives at least $\frac{1}{2}$ to f and $w'(f) \geq -2+4\times\frac{1}{2}=0$.

Case 3. $d(f)=5$.

We see that $w(f)=d(f)-6=-1$. If $n_{6^+}(f) \geq 1$, then by (R3) or (R4), each 6^+-vertex in $b(f)$ gives at least 1 to f. Hence, $w'(f)=-1+1\times 1=0$. So assume that $n_{6^+}(f)=0$.

This implies that $\delta(f) \geq 5$ and $n_5(f) = 5$. Then by (R2), each vertices in $b(f)$ gives at least $\frac{1}{2}$ to f and $w'(f) \geq -1 + 5 \times \frac{1}{2} = \frac{3}{2}$.

Case 4. $d(f) \geq 6$.

It is obvious that $w(f) = d(f) - 6 \geq 0$.

Let $v \in V(H)$. By Claim 1, we may assume that $d_H(v) \geq 3$.

If $d_H(v) = 3$, then $w(v) = 2d_H(v) - 6 = 0$.

Assume that $d_H(v) = 4$, then $w(v) = 2d_H(v) - 6 = 2$. If $n_{7^-}(v) = 1$, then $w'(v) \geq 2 - 2 \times \frac{4}{5} - 2 \times \frac{1}{5} = 0$ by (R1). Otherwise, $n_{7^-}(v) = 0$ and $w'(v) \geq 2 - 4 \times \frac{1}{2} = 0$.

Assume that $d_H(v) = 5$, then $w(v) = 2d_H(v) - 6 = 4$. If $n_4(v) \geq 1$, then $w'(v) \geq 4 - 5 \times \frac{4}{5} = 0$ by (R2). Otherwise, $n_4(v) = 0$. Let $w_1, w_2, \ldots, w_5$ be the neighbors of v in a cyclic order. We need to consider three subcases:

Case 1. $n_5(v) \geq 1$ with $d_H(w_1) = 5$.

Since G contains no $(A_{4.2})$, $n_6(v) \leq 2$. This leads to the following three possibilities:

(1.1) If $n_5(v) = 2$, then $n_{9^+}(v) = 3$ and $n'_{9^+}(v) = 3$ as G contains no $(A_{4.2})$. By symmetry, we may assume that $d_H(w_2) = 5$ or $d_H(w_3) = 5$. Thus, by (R2), $w'(f) \geq 4 - \max\{(\frac{7}{5} + 2 \times \frac{11}{12} + 2 \times \frac{1}{3}), (4 \times \frac{11}{12} + \frac{1}{3})\} = 4 - \max\{3\frac{9}{10}, 4\} = 0$.

(1.2) If $n_5(v) = n_6(v) = 1$, then $n_{8^+}(v) = 3$ and $n'_{8^+}(v) = 3$ as G contains no $(A_{4.2})$. By symmetry, we may assume that $d_H(w_2) = 6$ or $d_H(w_3) = 6$. Thus, by (R2), $w'(f) \geq 4 - \max\{(\frac{6}{5} + 1 + \frac{3}{4} + 2 \times \frac{1}{2}), (2 \times 1 + 2 \times \frac{3}{4} + \frac{1}{2})\} = 4 - \max\{3\frac{19}{20}, 4\} = 0$.

(1.3) Assume that $n_5(v) = 1$ and $n_{7^+}(v) = 4$. If $d_H(w_2) = d_H(w_5) = 7$, then by (R2), $w'(f) \geq 4 - (2 \times \frac{13}{14} + 3 \times \frac{5}{7}) = 0$. If one of w_2 and w_5 is of degree 7 in H, then by (R2), $w'(f) \geq 4 - (\frac{13}{14} + 1 + 2 \times \frac{5}{7} + \frac{9}{14}) = 0$. Otherwise, $\min\{d_H(w_2), d_H(w_5)\} \geq 8$, then by (R2), $w'(f) \geq 4 - (2 \times 1 + 2 \times \frac{9}{14} + \frac{5}{7}) = 0$.

Case 2. $n_5(v) = 0$ and $n_6(v) \geq 1$ with $d_H(w_1) = 6$.

Since G contains no $(A_{4.2})$, $n_6(v) \leq 2$. If $n_6(v) = 1$ and $n_{7^+}(v) = 4$, then by (R2), $w'(f) \geq 4 - (2 \times \frac{6}{7} + 3 \times \frac{5}{7}) = \frac{1}{7}$. Otherwise, $n_6(v) = 2$ and $n_{7^+}(v) = 3$. Since G contains no $(A_{4.2})$, $n_{8^+}(v) \geq 1$, or $n_7(v) = 3$ with $uv_5 \notin E(H)$ and $d_H(w_2) = 6$ or $d_H(w_3) = 6$. If $n_7(v) = 3$, then by (R2), v sends at most $\frac{1}{2}$ to the face whose boundary contains w_5, v, w_1. It follows that $w'(f) \geq 4 - (\frac{1}{2} + 1 + \frac{6}{7} + 2 \times \frac{5}{7}) = \frac{3}{14}$ if $d_H(w_2) = 6$, or $w'(f) \geq 4 - (\frac{1}{2} + 3 \times \frac{6}{7} + \frac{5}{7}) = \frac{3}{14}$ if $d_H(w_3) = 6$. Otherwise, $n_{8^+}(v) \geq 1$ and we need to consider the following subcases by symmetry.

- If $d_H(w_2) = 6$ and $d_H(w_3) \geq 8$, then by (R2), $w'(f) \geq 4 - (1 + \frac{6}{7} + \frac{5}{7} + \frac{3}{4} + \frac{9}{14}) = \frac{1}{28}$.
- If $d_H(w_2) = 6$ and $d_H(w_4) \geq 8$, then by (R2), $w'(f) \geq 4 - (1 + 2 \times \frac{6}{7} + 2 \times \frac{9}{14}) = 0$.
- If $d_H(w_3) = 6$ and $d_H(w_2) \geq 8$, then by (R2), $w'(f) \geq 4 - (2 \times \frac{6}{7} + 2 \times \frac{3}{4} + \frac{5}{7}) = \frac{1}{14}$.
- If $d_H(w_3) = 6$ and $d_H(w_4) \geq 8$, then by (R2), $w'(f) \geq 4 - (3 \times \frac{6}{7} + \frac{3}{4} + \frac{9}{14}) = \frac{1}{28}$.

Case 3. $n_5(v) = n_6(v) = 0$ and $n_{7^+}(v) = 5$. Then, by (R2), $w'(f) \geq 4 - 5 \times \frac{5}{7} = \frac{3}{7}$.

If $6 \leq d_H(v) \leq 9$, then by (R3), $w'(f) = 2d_H(v) - 6 - d_H(v) \times \frac{2d_H(v)-6}{d_H(v)} = 0$.

If $d_H(v) \geq 12$, then by (R4), $w'(f) \geq 2d_H(v) - 6 - d_H(v) \times \frac{3}{2} = \frac{1}{2}d_H(v) - 6 \geq \frac{1}{2} \times 12 - 6 = 0$.

Assume that $d_H(v) = 10$, then $w(f) = 2d_H(v) - 6 = 14$. Let t the number of faces f incident to v with $d(f) \geq 4$, or $d(f) = 3$ and $\delta(f) \geq 6$. If $t \geq 2$, then by (R4), $w'(f) \geq 14 - 8 \times \frac{3}{2} - 2 \times 1 = 0$. Otherwise, $t \leq 1$. Note that if $n_{5^-}(v) \leq 4$, then $t \geq 2$. Recall that if $d(v) \neq 10$, then $n_{5^-}(v) \leq 2 \leq 4$ by Claim 3. Hence, it suffice to assume that $d(v) = 10$ with $v_1, v_2, \ldots, v_{10}$ as the neighbors of v in a clockwise order, $n_{5^-}(v) \geq 5$ and $m_{4^+}(v) \leq t \leq 1$. If v is not incident to a bad 3-face, then $w'(f) = 14 - 10 \times \frac{7}{5} = 0$ by (R4). Otherwise, $n_3(v) \geq 1$ and v is incident to some bad 3-faces. Since G contains no $(A_{3.3})$, there is no 3-vertex w such that the faces f_1, f_2 which is incident to v, w, are both 3-faces. By the previous discussion, we may assume that $d_H(v_1) = 3$, $v_1v_2 \in E(H)$, $f_{10} = [\cdots v_{10}vv_1 \cdots]$ is a 4^+-face and the other faces incident to v are all of degree 3. Let $m'_3(v)$ be the number of bad 3-faces incident to v. Clearly, $1 \leq m'_3(v) \leq 2$, and $m'_3(v) = 2$ if and only if $d_H(v_{10}) = 3$. If $m'_3(v) = 2$, then $w'(f) \geq 14 - (2 \times \frac{3}{2} + 1 + 7 \times \frac{7}{5}) = \frac{1}{5}$ by (R4). Otherwise, $m'_3(v) = 1$ and hence $w'(f) \geq 14 - (1 \times \frac{3}{2} + 1 + 8 \times \frac{7}{5}) = \frac{3}{10}$.

Finally assume that $d_H(v) = 11$, then $w(f) = 2d_H(v) - 6 = 16$. If $n_3(v) \leq 3$, then the number of bad 3-faces incident to v is at most 6. Hence, by (R4), $w'(f) \geq 16 - 6 \times \frac{3}{2} - 5 \times \frac{7}{5} = 0$. Note that if $d(v) \neq 11$, then $n_3(v) \leq n_{5^-}(v) \leq 3$ by Claim 3. Now we assume that $d(v) = 11$ with $w_1, w_2, \ldots, w_{11}$ as the neighbors of v in a clockwise order, $n_3(v) \geq 4$ and v is incident to at least 7 bad 3-faces. If v is incident to a 4^+-face, or to a 3-face f with $\delta(f) \geq 6$, then $w'(f) = 16 - 10 \times \frac{3}{2} - 1 = 0$. Otherwise, $m_3(v) = 11$ and $\delta(f) \leq 5$ for each f incident to v, i.e., for $1 \leq i \leq 11$ (where $w_{12} = w_1$), $\min\{d_H(w_i), d_H(w_{i+1})\} \leq 5$.

Note that if $\min\{d_H(w_i), d_H(w_{i+1})\} = 3$, then $\max\{d_H(w_i), d_H(w_{i+1})\} \geq 10$ by Claim 4 for each $1 \leq i \leq 11$ (where $w_{12} = w_1$). Thus, $n_3(v) \leq 5$. If $n_3(v) = 5$, then we may assume that $d_H(w_i) = 3$ for $i = 3, 5, 7, 9, 11$ and $d_H(w_j) \geq 10$ for $j = 1, 2, 4, 6, 8, 10$. Obviously, $f_1 = [w_1vw_2]$ is of degree 3 with $\delta(f) \geq 10$, a contradiction. Otherwise, $n_3(v) = 4$. Further, for $1 \leq i \leq 11$ (where $w_{12} = w_1$, $w_{13} = w_2$, $w_{14} = w_3$), if $d_H(w_i) = d_H(w_{i+3}) = 3$, then $d_H(w_{i+1}) \geq 10, d_H(w_{i+2}) \geq 10$ by Claim 4 and $f_{i+1} = [w_{i+1}vw_{i+2}]$ is of degree 3 with $\delta(f_{i+1}) \geq 10$, a contradiction. Hence, it suffice to assume that $d_H(w_i) = 3$ for $i = 4, 6, 8, 10$; $d_H(w_j) \geq 10$ for $j = 3, 5, 7, 9, 11$ and $4 \leq d_H(w_1), d_H(w_2) \leq 5$.

Since G contains no $(A_{4.1})$, there is no 4-vertex adjacent to a 4-vertex and 11-vertex at the same time and $\max\{d_H(w_1), d_H(w_2)\} = 5$. Without loss of generality, assume that $d_H(w_1) = 5$. If $d_H(w_2) = 4$, then $n_4(w_1) \geq 1$ and $n_{7^-}(w_2) = 1$. By (R4), $w'(f) = 16 - 8 \times \frac{3}{2} - 2 \times \frac{7}{5} - \frac{11}{10} = \frac{1}{10}$. Otherwise, $d_H(w_2) = 5$. If $n_4(w_1) \geq 1$ and $n_4(w_2) \geq 1$, then By (R4), $w'(f) = 16 - 8 \times \frac{3}{2} - 2 \times \frac{11}{10} - \frac{7}{5} = \frac{2}{5}$. If $n_4(w_1) = n_4(w_2) = 0$, then By (R4), $w'(f) = 16 - 8 \times \frac{3}{2} - 3 \times \frac{4}{3} = 0$. Otherwise, by (R4), $w'(f) = 16 - 8 \times \frac{3}{2} - \frac{7}{5} - \frac{4}{3} - \frac{11}{10} = \frac{1}{6}$. □

3 Acyclic chromatic indices

In this section, we discuss the acyclic chromatic indices of planar graphs. The proof of the main result requires the following lemmas.

Lemma 2 ([12]) *If G is a graph with $\Delta \leq 4$, then $a'(G) \leq 7$.*

Assume that c is a partial acyclic edge k-coloring of a graph G using the color set $C = \{1, 2, \ldots, k\}$. For a vertex $v \in V(G)$, we use $C(v)$ to denote the set of colors assigned to edges incident to v under c. If the edges of a cycle are alternatively colored with colors i and j, then we call such cycle an *(i, j)-cycle*. If the edges of a path $ux \ldots v$ are alternatively colored with colors i and j, then we call such path an *$(i, j)_{(u,v)}$-path*.

Lemma 3 *If $P = uv_1v_2 \ldots v_kv_{k+1}$ is an $(i, j)_{(u,v_{k+1})}$-path in G with $c(uv_1) = i$, $j \notin C(u)$ and $w \notin V(P)$, then there is no $(i, j)_{(u,w)}$-path in G.*

Proof. Otherwise, assume that there is an $(i, j)_{(u,w)}$-path $Q = uw_1w_2 \ldots w_m$ in G, where $w = w_m$ and $m \geq 1$. Note that some w_i may be identical to some v_j, $1 \leq i \leq k + 1$ and $1 \leq j \leq m - 1$. Since $j \notin C(u)$ and $w \notin V(P)$, it is easy to see that there exist a vertex v_i and some vertex w_i such that v_iv_{i+1} and v_iw_i have same color i or j, which contradicts the fact that the edge coloring considered is proper. $\square$

Theorem 1 *If G is a planar graph, then $a'(G) \leq \Delta + 7$.*

Proof. The proof is proceeded by induction on the edge number $|E(G)|$. If $|E(G)| \leq \Delta+7$, then G is obviously acyclically edge $(\Delta+7)$-colorable. Assume that G is a planar graph with $|E(G)| \geq \Delta + 8$. If $\Delta \leq 4$, then $a'(G) \leq \Delta + 3$ by Lemma 2. Thus, we may assume that $\Delta \geq 5$ and G is 2-connected. By Lemma 1, G contains one of the configurations (A_1)-(A_4). In the following, we deal with each of the configurations (A_1)-(A_4).

(A$_1$) There is a path uvw such that $d(v) = 2$ and $d(u) \leq 9$.

Let $v, u_1, u_2, \ldots, u_{d(u)-1}$ be the neighbors of u and let $H = G - uv$. Then H is a planar graph and $\Delta(H) \geq \Delta - 1 \geq 4$. By the induction assumption or Lemma 2, H has an acyclic edge $(\Delta + 7)$-coloring c using the color set $C = \{1, 2, \ldots, \Delta + 7\}$.

If $c(vw) \notin C(u)$, then we color uv with a color in $C\backslash(C(u) \cup C(v))$. Otherwise, $c(vw) \in C(u)$ and assume that $c(vw) = c(uu_1) = 1$, $c(uu_i) = i$, $i = 2, 3, \ldots, d(u)-1$. If H contains no $(1, a)_{(w,u_1)}$-path for some color $a \in C\backslash(C(u) \cup C(v))$, we color uv with a. Otherwise, assume that H contains a $(1, i)_{(w,u_1)}$-path for any $i \in C\backslash(C(u) \cup C(v))$. This implies that $d(u) = 9$ and $C(w) = C(u_1) = (C\backslash C(v)) \cup \{1\} = C\backslash\{2, 3, 4, 5, 6, 7, 8\}$. First, we recolor vw with 2. Similarly to the previous argument, we may assume that H contains a $(2, i)_{(u_2,w)}$-path for any $i \in C\backslash(C(u) \cup C(v))$ and $C(u_2) = C\backslash\{1, 3, 4, 5, 6, 7, 8\}$. So we switch the colors of uu_1 and uu_2, then color uv with 9. Clearly, no bichromatic cycles are produced in G under the constructed coloring by the previous assumption.

(A$_2$) There is a vertex u with $n_2(u) \geq 1$ and $n_{8^-}(u) \geq d(u) - 8$. Suppose that $u_1, u_2, \ldots, u_{d(u)-1}$, v are the neighbors of u such that $d(u_1) \geq d(u_2) \geq \cdots \geq d(u_{d(u)-1}) \geq d(v) = 2$. Let w be the neighbor of v different from u. For $1 \leq i \leq d(u) - 1$, if $d(u_i) = 2$, then we use x_i to denote the neighbor of u_i different from u. Then at least one of the configurations $(A_{2.1})$-$(A_{2.2})$ holds.

Without loss of generality, assume that $c(uu_i) = i$ for $1 \leq i \leq d(u)-1$ and let $S_{i^-} = \{c(uu_j)|d(u_j) \leq i\}$. Since $d(u) \leq \Delta$, $|C\backslash C(u)| \geq 8$. If $c(vw) \notin C(u)$, then we color uv with a color in $C\backslash(C(u) \cup C(v))$. Otherwise, assume that $c(vw) \in C(u)$ and H contains a $(c(vw), i)_{(u,w)}$-path for any $i \in C\backslash C(u)$, $(C\backslash C(u)) \subseteq C(w)$. If $c(vw) = c(uu_i) \in S_{8^-}$, we color uv with a color in $(C\backslash C(u))\backslash C(u_i)$. Otherwise, $c(vw) \in C(u)\backslash S_{8^-}$. Further, we may assume that $S_{8^-} \subseteq C(w)$. Otherwise, we recolor vw with a color in $(S_{8^-})\backslash C(w)$ and reduce the proof to the case in which $c(vw) \in S_{8^-}$. So we assume that:

$(*_{2.1})$ $c(vw) \in C(u)\backslash S_{8^-}$ and $S_{8^-} \cup (C\backslash C(u)) \subseteq C(w)\backslash\{c(vw)\}$.

$(\mathbf{A}_{2.1})$ $n_{8^-}(u) \geq d(u) - 7$.

By $(*_{2.1})$, $(C\backslash\{c(uu_1), c(uu_2), \cdots, c(uu_7)\}) \subseteq C(w)\backslash\{c(vw)\}$ and $d(w) = |C(w)| \geq \Delta+1$.

$(\mathbf{A}_{2.2})$ $n_{8^-}(u) = d(u) - 8$, and $n_2(u) \geq d(u) - 9$.

Without loss of generality, we assume that $d(u_9) = 8$. By $(*_{2.1})$, we assume that $d(u_i) \geq 9$, $i = 1, 2, \cdots, 8$, $c(vw) = 1$ and $C(w) = C\backslash\{2, 3, \ldots, 8\}$. Further, we assume that H contains a $(1, i)_{(u_1,w)}$-path for any $i \in C\backslash C(u)$. If H contains no $(i, j)_{(u_i,w)}$-path for some $j \in C\backslash C(u)$ and some $i \in \{2, 3, \cdots, 8\}$, we recolor vw with i and color uv with j. Otherwise, we may assume that:

$(*_{2.2})$ For any $i \in \{1, 2, \cdots, 8\}$ and $j \in C\backslash C(u)$, H contains an $(i, j)_{(u_i,w)}$-path and $(C\backslash C(u)) \subseteq C(u_i)$.

Assume that H contains no $(i, 9)_{(u_i,w)}$-path for some color $i \in \{1, 2, \ldots, 8\}$. Let $C(u_9) = \{9, a_1, a_2, \ldots, a_7\}$. If $\{a_1, a_2, \ldots, a_7\} \cap S_2 = \emptyset$, then we recolor uu_9 with a color in $C\backslash(C(u) \cup C(u_9))$. Otherwise, assume that $\{a_1, a_2, \ldots, a_7\} \cap S_2 \neq \emptyset$. Let $I = \{i|i \in \{10, \ldots, d(u)-1\}$ such that $c(uu_i) \in \{a_1, a_2, \ldots, a_7\}\}$. Then we recolor uu_9 with a color in $C\backslash(C(u) \cup C(u_9) \cup \{C(u_i)|i \in I\})$, vw with i if $i \neq 1$ and color uv with 9. So we may assume that:

$(*_{2.3})$ H contains a $(i, 9)_{(u_i,w)}$-path for any $i \in \{1, 2, \ldots, 8\}$.

Assume that H contains no $(i, j)_{(u_i,w)}$-path for some $i \in \{1, 2, \ldots, 8\}$ and some $j = c(uu_j) \in S_2$. Let $C(u_j) = \{j, a\}$. If $a \notin S_2 \cup \{9\}$, we recolor uu_j with a color in $C\backslash(C(u) \cup C(u_j))$; If $a = \{c(uu_k)\} \in S_2 \cup \{9\}$, we recolor uu_j with a color in $C\backslash(C(u) \cup C(u_j) \cup C(u_k))$. Recolor vw with i if $i \neq 1$ and color uv with j. So we may assume that:

$(*_{2.4})$ H contains an $(i, j)_{(u_i,w)}$-path for any $i \in \{1, 2, \ldots, 8\}$ and any $j \in S_2$.

By $(*_{2.2})$, $(*_{2.3})$, and $(*_{2.4})$, we may assume that:

$(*_{2.5})$ For any $j \in C\backslash\{1, 2, \ldots, 8\}$, H contains an $(i, j)_{(u_i,w)}$-path, and $C(u_i)\backslash \{i\} = C(w) \backslash\{c(vw)\} = C\backslash \{1, 2, \ldots, 8\}$, $i = 1, 2, \ldots, 8$.

Claim 8 *There exist $i_0, j_0 \in \{1, 2, \ldots, 8\}$ such that H contains neither an $(i_0, 9)_{(u_{j_0},u_9)}$-path nor a $(j_0, 9)_{(u_{i_0},u_9)}$-path.*

Proof. Since $d(u_9) = 8$, $\{1, 2, \dots, 8\}\backslash C(u_9) \neq \emptyset$. Assume that $1 \notin C(u_9)$. If H contains no $(2, 9)_{(u_1,u_9)}$-path, $\{i_0, j_0\} = \{1, 2\}$. If H contains no $(3, 9)_{(u_1,u_9)}$-path, $\{i_0, j_0\} = \{1, 3\}$. Otherwise, H contains a $(2, 9)_{(u_1,u_9)}$-path and a $(3, 9)_{(u_1,u_9)}$-path. It follows that H contains neither a $(2, 9)_{(u_3,u_9)}$-path nor a $(3, 9)_{(u_2,u_9)}$-path, thus $\{i_0, j_0\} = \{2, 3\}$. □

By symmetry, assume that $\{i_0, j_0\} = \{1, 2\}$. We switch the colors of uu_1 and uu_2. If no bichromatic cycles are produced, we color uv with some color in $C\backslash C(u)$ and we are done. Otherwise, by the previous assumption, it is easy to conclude that the bichromatic cycle in G must be a $(1, i)_{(u_2,u_i)}$-cycle or a $(2, j)_{(u_1,u_j)}$-cycle for some $i, j \in S_2$. Let

$$T_1 = \{u_i|\ G \text{ contains a } (1, i)_{(u_2,u_i)}\text{-cycle}\},$$

$$T_2 = \{u_i|\ G \text{ contains a } (2, i)_{(u_1,u_i)}\text{-cycle}\}.$$

If $|T_1| \geq 2k$, $k \geq 1$, assuming that $u_{i_1}, u_{i_2} \in T_1$, then we switch the colors of uu_{i_1} and uu_{i_2}, so that G contains neither $(1, i_1)_{(u_2,u_{i_1})}$-cycle nor $(1, i_2)_{(u_2,u_{i_2})}$-cycle, and no other new bichromatic cycles are produced. Since $\min\{|T_1|, |T_2|\} \geq 1$, we assume that $|T_1| = 1$, $|T_2| \leq 1$. Without loss of generality, assume that $u_{10} \in T_1$, $c(u_{10}x_{10}) = 1$, and $10 \in C(x_{10})$.

Since $d(x_{10}) \leq \Delta$, $(C\backslash\{2\})\backslash C(x_{10}) \neq \emptyset$, assume that $p \in (C\backslash\{2\})\backslash C(x_{10})$ and recolor $u_{10}x_{10}$ with p. If $p \in (C\backslash C(u)) \cup \{3, 4, \dots, 8\}$, then let $S'_2 = S_2$ and $C'(u) = C(u)$. If $p = c(uu_i) \in S_2 \cup \{9\}$, recolor uu_{10} with a color $j_1 \in (C\backslash C(u))\backslash C(u_i)$ and let $S'_2 = (S_2\backslash\{10\}) \cup \{j_1\}$, $C'(u) = (C(u)\backslash\{10\}) \cup \{j_1\}$. If $|T_2| = 0$, then we color uv with 10. Otherwise, $|T_2| = 1$, $u_{11} \in T_2$ and $c(u_{11}x_{11}) = 11$. Since $d(x_{11}) \leq \Delta$, $(C\backslash\{1\})\backslash C(x_{11}) \neq \emptyset$, assume that $q \in (C\backslash\{1\})\backslash C(x_5)$ and recolor $u_{11}x_{11}$ with q. If $q \in (C\backslash C'(u)) \cup \{3, 4, \dots, 8\}$, then let $C''(u) = C'(u)$. If $q = c(uu_i) \in S'_2 \cup \{9\}$, we recolor uu_{11} with a color $j_2 \in (C\backslash C'(u))\backslash C(u_i)$ and $C''(u) = (C'(u)\backslash\{11\}) \cup \{j_2\}$. Then, we color uv with a color in $C\backslash C''(u)$.

($\mathbf{A}_3$) There is a 3-vertex u adjacent to a vertex v such that one of $(A_{3.1})$-$(A_{3.3})$ holds:

Let $v_1, v_2, \dots, v_{d(v)}$ be the neighbors of v, where $v_8 = u_2$ in $(A_{3.2})$, ($v_8 = u_1$, $v_9 = u_2$ in $(A_{3.3})$) and $v_{d(v)} = u$. Let $H = G - uv$. Then H is a planar graph and $\Delta(H) \geq \Delta - 1 \geq 4$. By the induction assumption or Lemma 2, H has an acyclic edge $(\Delta + 7)$-coloring c using the color set $C = \{1, 2, \dots, \Delta + 7\}$. Note that $|C\backslash C(v)| \geq 8$. If $C(u) \cap C(v) = \emptyset$, then color uv with a color in $C\backslash(C(u) \cup C(v))$. Otherwise, $C(u) \cap C(v) \neq \emptyset$, i.e., $1 \leq |C(u) \cap C(v)| \leq 2$ and $c(uu_i) = i$, $i = 1, 2$.

If, for any $i \in C(u) \cap C(v)$, H contains no $(i, j)_{(u,v)}$-path for some $j \in C\backslash(C(u) \cup C(v))$, then we can color uv with j. Otherwise, we assume that:

$(*_{3.1})$ For any $j \in C\backslash(C(u) \cup C(v))$, H contains an $(i, j)_{(u,v)}$-path for some $i \in C(u) \cap C(v)$.

Next we need to consider the following two subcases.

(I) $|C(u) \cap C(v)| = 1$.

($\mathbf{A}_{3.1}$) $d(v) \leq 8$.

Assume that $C(v) = \{1, 2, \dots, d(v)\}\backslash\{2\}$, and $c(uu_1) = c(vv_1) = 1$. By $(*_{3.1})$, $d(v) = 8$, $C(u_1) = C(v_1) = C\backslash\{2, 3, \dots, 8\}$. We recolor uu_1 with 3, vv_1 with 2, and color uv with 1. Clearly, no bichromatic cycles are produced in G under the constructed coloring.

($\mathbf{A}_{3.2}$) $d(v) = 9$ and $uu_2, vu_2 \in E(G)$.

(3.2.1) $C(v) = \{1, 2, \ldots, d(v)\}\backslash\{2\}$.

Assume that $c(vu_2) = 1$. By $(*_{3.1})$, $d(u_2) = \Delta$, $C(v) \cup C(u_2) = C$, $C(v) \cap C(u_2) = \emptyset$ and $C(u_2)\backslash\{2\} \subseteq C(u_1)$. Since $d(u_1) \le \Delta$, there exists $a_1 \in C(v)\backslash C(u_1)$. We recolor uu_2 with a_1 and color uv with 2.

Assume that $c(vv_i) = i$, $i \in \{1, 2, \ldots, 7\}\backslash\{2\}$, and $c(vu_2) = 8$, $c(vv_2) = 9$. By $(*_{3.1})$, $C\backslash(2, 3, \ldots, 9) \subseteq C(u_1)$. Since $d(u_1) \le \Delta$, $|\{2, 3, \ldots, 9\} \cap C(u_1)| \le 1$. If $8 \notin C(u_1)$, we recolor uu_1 with 8 and no bichromatic cycle is produced since $2 \notin C(v)$, then the proof is similar to the case $c(vu_2) = c(uu_1) = 1$. Otherwise, $C(u_1) = (C\backslash(C(u) \cup C(v))) \cup \{1, 8\}$. If there exists $a_2 \in (C\backslash(C(u) \cup C(v)))\backslash C(u_2)$, we recolor uu_2 with a_2 and color uv with 2. Otherwise, $C(u_2) = (C\backslash(C(u) \cup C(v))) \cup \{2, 8\}$. Then, we recolor uu_2 with 1, uu_1 with 3 and color uv with 2.

(3.2.2) $C(v) = \{1, 2, \ldots, d(v)\}\backslash\{1\}$.

Assume that $c(vv_i) = i$, $i \in \{1, 2, \ldots, 7\}\backslash\{1\}$, and $c(vu_2) = 8$, $c(vv_1) = 9$. By $(*_{3.1})$, $d(u_2) = \Delta$, $C(v) \cup C(u_2) = C$, $C(v) \cap C(u_2) = \emptyset$. Then, we recolor uu_1 with a color in $C\backslash(C(u_1) \cup \{2, 8\})$ and color uv with 1.

($A_{3.3}$) $d(v) = 10$, $n_{5^-}(v) \ge 5$ and $uu_1, vu_1, uu_2, vu_2 \in E(G)$.

Without loss of generality assume that $C(v) = \{1, 2, \ldots, d(v)\}\backslash\{2\}$ and $c(vv_i) = i$, $i \in \{3, 4, \ldots, 7\}$, $c(vu_1) = 8$, $c(vv_2) = 10$.

(3.3.1) $c(vu_2) = 1$ and $c(vv_1) = 9$.

By $(*_{3.1})$, $(C\backslash(C(u) \cup C(v))) \subseteq C(u_1) \cap C(u_2)$. Since $d(u_1), d(u_2) \le \Delta$, $|\{3, 4, \ldots, 7, 9, 10\} \cap C(u_1)| \le 1$, and $|\{3, 4, \ldots, 7, 9, 10\} \cap C(u_2)| \le 1$. Thus, there exists $a_3 \in \{3, 4, \ldots, 7, 9, 10\}\backslash (C(u_1) \cup C(u_2))$. Then, we recolor uu_2 with a_3 and color uv with 2.

(3.3.2) $c(vu_2) = 9$ and $c(vv_1) = 1$.

By $(*_{3.1})$, $(C\backslash(C(u) \cup C(v))) \subseteq C(u_1) \cap C(v_1)$. If $9 \notin C(u_1)$, we recolor u_1u with 9 and the proof is similar to (3.3.1). Otherwise, $9 \in C(u_1)$. Since $d(u_1) \le \Delta$, $d(u_1) = \Delta$, and $C(u_1) = (C\backslash(C(u) \cup C(v))) \cup \{1, 8, 9\}$. Obviously, $2 \notin C(u_1)$. If there exists $a_4 \in (C\backslash C(u) \cup C(v))\backslash C(u_2)$, recolor uu_2 with a_4 and color uv with 2. By $(*_{3.1})$ and Lemma 3, no bichromatic cycles are produced. Otherwise, $(C\backslash C(u) \cup C(v)) \subseteq C(u_2)$ and there exists $a_5 \in \{3, 4, \ldots, 7, 10\}\backslash C(u_2)$. Then, we recolor uu_2 with a_5 and color uv with 2.

(II) $|C(u) \cap C(v)| = 2$.

If there exists $a_6 \in C\backslash(C(u) \cup C(v) \cup C(u_{i_0}))$ for some $i_0 \in \{1, 2\}$, then we recolor uu_{i_0} with a_6 and reduce the proof to the Case I. Otherwise, we assume that:

$(*_{3.2})$ For each $i \in \{1, 2\}$, $C\backslash(C(u) \cup C(v) \cup C(u_i)) = \emptyset$, i.e., $(C(u) \cup C(v) \cup C(u_i)) = C$.

Note that in $(A_{3.1})$ and $(A_{3.2})$, $C\backslash(C(u) \cup C(v) \cup C(u_2)) \neq \emptyset$. So we assume that $d(v) = 10$, $n_{5^-}(v) \ge 5$ and $uu_i, vu_i \in E(G)$, $i = 1, 2$. Further, assume that $C(v) = \{1, 2, \ldots, 9\}$ and $c(vv_i) = i$, $i = \{1, 2, \ldots, 9\}\backslash\{2, 8\}$.

- $c(vu_1) = 8$, $c(vv_2) = 2$.

By $(*_{3.2})$, $d(u_i) = \Delta$, $i = 1,2$ and $C(u_1) = (C\backslash\ \{1,2,\ldots,\ 9\}) \cup\ \{1,8\}$, $C(u_2) = (C\backslash\{1,2,\ldots,9\}) \cup \{2,9\}$. Since $n_{5^-}(v) \geq 5$, there exists $j = c(vv_j), k = c(vv_k)$ and $d(v_j), d(v_k) \leq 5$. Without loss of generality, assume that $j = 1$ and $k = 2$ (Otherwise, we can recolor uu_1 with j, uu_2 with k if $\{j,k\} \neq \{1,2\}$ and the proof is similar. Clearly, no bichromatic cycles are produced and $C(u) \cap C(v) = \{j,k\}$.) By $(*_{3.1})$, for any $j \in C\backslash(C(u) \cup C(v))$, H contains an $(i,j)_{(u,v)}$-path for some color $i \in \{1,2\}$. Hence, $\Delta = 10$, $C(u_1) = \{1,10,11,12,13\}$ and $C(u_2) = \{2,14,15,16,17\}$. Then, switch the colors of uu_1 and uu_2, and color uv with 10. By $(*_{3.1})$ and Lemma 3, no bichromatic cycles are produced.

- $c(vu_1) = 2$, $c(vv_2) = 8$.

By $(*_{3.2})$, $d(u_i) = \Delta$, $i = 1,2$ and $C(u_1) = (C\backslash\ \{1,2,\ldots,\ 9\}) \cup \{1,2\}$, $C(u_2) = (C\backslash\{1,2,\ldots,9\}) \cup \{2,9\}$. We recolor uu_2 with 8 and the proof is similar to the case $c(vu_1) = 8$ and $c(uu_2) = 2$.

($\mathbf{A}_4$) There is a vertex v adjacent to $u, v_2, \ldots, v_{d(v)}$ with $d(u) \leq d(v_2) \leq \ldots \leq d(v_{d(v)})$ such that one of $(A_{4.1})$-$(A_{4.2})$ holds.

Let $u_1, u_2, \ldots, u_{d(u)-1} \neq v$ be the other neighbors of u. Let $H = G - uv$. Then H is a planar graph and $\Delta(H) \geq \Delta - 1 \geq 4$. By the induction assumption or Lemma 2, H has an acyclic edge $(\Delta + 7)$-coloring c using the color set $C = \{1, 2, \ldots, \Delta + 7\}$ with $c(uu_i) = i$, $i = 1, 2, \ldots, u_{d(u)-1}$. Note that $|C\backslash C(u)| \geq 8$, $|C(v)| \leq d(v) \leq 5$ and $C\backslash(C(u) \cup C(v)) \neq \emptyset$. If $C(u) \cap C(v) = \emptyset$, then we color uv with a color in $C\backslash(C(u) \cup C(v))$. Otherwise, $C(u) \cap C(v) \neq \emptyset$.

If, for any $i \in C(u) \cap C(v)$, H contains no $(i,j)_{(u,v)}$-path for some $j \in C\backslash(C(u) \cup C(v))$, then we can color uv with j. Otherwise, we assume that:

$(*_{4.1})$ For any $j \in C\backslash(C(u) \cup C(v))$, H contains an $(i,j)_{(u,v)}$-path for some $i \in C(u) \cap C(v)$.

Next we need to consider the following four subcases.

Case 1 $|C(u) \cap C(v)| = 1$ and $1 = c(uu_1) \in C(v)$.

By $(*_{4.1})$, we only need to consider two cases: (i) $d(u) = 7$, $d(v) = 4$ and $C(v) = \{1,7,8\}$ with $c(w_1) = 1$, $w_1, w_2 \in \{v_1, v_2, v_3\}$; (ii) $d(u) = 6$, $d(v) = 5$ and $C(v) = \{1,6,7,8\}$ with $c(w_1) = 1$, $w_1, w_2, w_3 \in \{v_1, v_2, v_3, v_4\}$. By $(*_{4.1})$, we can assume that $C(u_1) = C(w_1) = (C\backslash\{1,2,\ldots,8\}) \cup \{1\}$. Then, we recolor uu_1 with 7, vw_1 with 2 and color uv with 1. Since $7, 8 \notin C(w_1)$ and $1 \notin C(u_1)$, no bichromatic cycles are produced.

Otherwise, $|C(u) \cap C(v)| \geq 2$ and $|C\backslash(C(u) \cup C(v))| = \Delta + 7 - |C(u) \cup C(v)| = \Delta + 7 - (|C(u)| + |C(v)| - |C(u) \cap C(v)|) = \Delta + 7 + |C(u) \cap C(v)| - (d(u) - 1 + d(v) - 1) = \Delta + 9 + |C(u) \cap C(v)| - (d(u) + d(v)) \geq \Delta + 9 + |C(u) \cap C(v)| - 11 \geq \Delta + 9 + 2 - 11 = \Delta$.

Let $\{w_1, w_2, \ldots, w_{d(v)-1}\} = \{v_2, v_3, \ldots, v_{d(v)}\}$. For any $1 \leq i \leq |C(u) \cap C(v)|$, let $c(vw_i) = i$, (i.e., $C(u) \cap C(v) = \{i | 1 \leq i \leq |C(u) \cap C(v)|\}$), $C_i = \{j \in C\backslash(C(u) \cup C(v)) |\ H$ contains an $(i,j)_{(u_i,v)}$-path$\}$ and $T_i = (C\backslash(C(u) \cup C(v)))\backslash C(w_i)$. In particular, $c(vw_1) = 1$, $c(vw_2) = 2$ and $T_1 = (C\backslash(C(u) \cup C(v)))\backslash C(w_1) = \{\beta_i | i = 1, 2, \ldots\}$, $T_2 = (C\backslash(C(u) \cup C(v)))\backslash C(w_2) = \{\alpha_i | i = 1, 2, \ldots\}$.

To complete the proof, we introduce a few symbols as defined in [5]. A multiset is a generalized set where a member can appear multiple times. If an element x appears t times in the multiset S, then we say that the multiplicity of x in S is t, and write $\mathrm{mult}_S(x) = t$. The cardinality of a finite multiset S, denoted by $\|S\|$, is defined as $\|S\| = \sum_{x\in S}\mathrm{mult}S(x) = t$. Let S_1 and S_2 be two multisets. The join of S_1 and S_2, denoted $S_1 \biguplus S_2$, is a multiset that have all the members of S_1 as well as S_2. For $x \in S_1 \biguplus S_2$, $\mathrm{mult}_{S_1 \biguplus S_2}(x) = \mathrm{mult}_{S_1}(x)+\mathrm{mult}_{S_2}(x)$. Clearly, $\|S_1 \biguplus S_2\| = \|S_1\| + \|S_2\|$.

Moreover, we define

$$S_v = \biguplus_{1\le i\le d(v)-1} (C(w_i)\backslash\{c(vw_i)\}).$$

Clearly, $\|S_v\| = \sum_{1\le i\le d(v)-1} d(w_i) - (d(v) - 1) = \sum_{2\le i\le d(v)} d(v_i) - (d(v) - 1)$ and for any $x \in C\backslash(C(u) \cup C(v))$, $\mathrm{mult}_{S_v}(x) \ge 1$ by $(*_{4.1})$. Further, for any $i \in C(u) \cap C(v)$, let

$$T'_i = \{x \in T_i|\mathrm{mult}_{S_v}(x) = 2\},$$

$$T_0 = \{x \in \biguplus_{i\in C(u)\cap C(v)} T_i|\mathrm{mult}_{S_v}(x) \ge 3\}.$$

Claim 9 *Let* $|C(u) \cap C(v)| = k + 1 \ge 2$. *If we can obtain an acyclic edge coloring* c *of* G *when* $|C(u) \cap C(v)| = k$, *then for any* $x \in C\backslash(C(u) \cup C(v))$, $\mathrm{mult}_{S_v}(x) \ge 2$ *and* $\|S_v\| \ge 2|C\backslash(C(u) \cup C(v))|$.

Proof. If not, then there exists $x \in C\backslash(C(u)\cup C(v))$ such that $\mathrm{mult}S_v(x) = 1$ and $x \in C(w_1)$. Obviously, $x \in C_1$ by $(*_{4.1})$ and the $(1, x)_{(u,v)}$-path cannot pass through w_2. We recolor vw_2 with x. Since $\mathrm{mult}S_v(x) = 1$, $x \notin C(w_i)$, $i \ne 1$ and no bichromatic cycles are produced. Then, we reduce the proof to the Case $|C(u) \cap C(v)| = k$. □

Case 2 $|C(u) \cap C(v)| = 2$ and $C(u) \cap C(v) = \{1, 2\}$.

By $(*_{4.1})$, for any $j \in C\backslash(C(u) \cup C(v))$, H contains an $(1, j)_{(u_1,v)}$-path or $(2, j)_{(u_2,v)}$-path and $T_1 \subseteq C_2 \subseteq C(w_2)$, $T_2 \subseteq C_1 \subseteq C(w_1)$, $(C\backslash(C(u) \cup C(v))) \subseteq C_1 \cup C_2$. And for any $i \in T_1$, $(2, i)_{(u_2,w_2)}$-path cannot path through w_1, while for any $j \in T_2$, $(1, j)_{(u_1,w_1)}$-path cannot pass through w_2. Further, by Claim 9, $T_1 \subseteq \bigcup_{3\le i\le d(v)-1} C(w_i)$ and $T_2 \subseteq \bigcup_{3\le i\le d(v)-1} C(w_i)$.

Claim 10 $\|S_v \cap C(u)\| \ge |C(u)| - 1 = d(u) - 2$.

Proof. If not, then there are at least two colors $x_1, x_2 \in C(u)\backslash S_v$. First, remove the colors of vw_1 and vw_2, then color vw_1 with x_1, vw_2 with x_2, i.e., $c(vw_1) = x_1$, $c(vw_2) = x_2$. Since $x_1, x_2 \notin S_v$, no bichromatic cycles are produced. Without loss of generality, we assume that $x_1 = 1$ and $x_2 = 2$. First, we switch the colors of vw_1 and vw_2. Next, if $C_1 \cap C_2 \ne \emptyset$, then we color uv with a color in $C_1 \cap C_2$. Otherwise, $C_1 \cap C_2 = \emptyset$. Since $d(u_1) \le \Delta$, $|C_1| + |C_2| = |C_1 \cup C_2| \ge |C\backslash(C(u) \cup C(v))| \ge \Delta$ and $1 \in C(u_1)$, there exits $b_1 \in C_2\backslash C(u_1)$. Then we color uv with b_1 and no bichromatic cycles are produced by $(*_{4.1})$ and Lemma 3. □

Claim 11 *For any* $i \in C\{1, 2\}$, $(C(v)\backslash\{1, 2\}) \cap C(w_i) \ne \emptyset$.

Proof. If not, $(C(v)\backslash\{1,2\})\cap C(w_{i_0})=\emptyset$ for some $i_0\in\{1,2\}$, then we recolor vw_{i_0} with a color in T_{i_0} and reduce the proof to the Case 1. $\square$

Recall that $|C\backslash(C(u)\cup C(v))|\geq\Delta$. It follows that $|T_i|\geq 2$ for any $i\in\{1,2\}$.

Case 2.1 $(A_{4.1})$ holds and $d(v)=4$.

Assume that $c(vw_3)=7$. By Claim 11, $7\in C(w_1)$ and $7\in C(w_2)$. By Claim 9 and Claim 10, for any $x\in C\backslash(C(u)\cup C(v))$, $\text{mult}_{S_v}(x)\geq 2$ and $\|S_v\cap C(u)\|\geq d(u)-2$. And recall that $7\in C(w_1)\cap C(w_2)$, we have $\|S_v\|\geq 2|C\backslash(C(u)\cup C(v))|+(|C(u)|-1)+|\{7\}|+|\{7\}|=2(\Delta+11-(d(u)+d(v)))+(d(u)-2)+2=2\Delta+14-d(u)$. However, $\|S_v\|=\sum_{2\leq i\leq d(v)}d(v_i)-(d(v)-1)=\sum_{2\leq i\leq d(v)}d(v_i)-3\leq 2\Delta+d(v_2)-3$. Clearly, $2\Delta+14-d(u)\leq 2\Delta+d(v_2)-3$, which implies that $d(u)+d(v_2)\geq 17$. Hence, $d(u)+d(v_2)=17$ and $\|S_v\cap C(u)\|=d(u)-2$. Assume that there exists $k\in C(u)\backslash S_v$. By $(*_{4.1})$, for any $i\in T_1$, $(2,i)_{(u,v)}$-path cannot pass through w_1 since $i\notin C(w_1)$. If $k=1$, recolor vw_3 with 1, vw_1 with β_1 and color uv with α_1. Similarly, $k\neq 2$. Otherwise, $k\in C(u)\backslash\{1,2\}$. Without loss of generality, assume that $k=3$. If H contains no $(3,\beta_1)_{(u,w_2)}$-path, we recolor vw_2 with 3 and color uv with β_1. Otherwise, recolor vw_3 with 3, vw_2 with α_1 and color uv with β_1. By $(*_{4.1})$ and Lemma 3, no bichromatic cycles are produced.

Case 2.2 $(A_{4.2})$ holds and $d(v)=5$.

Assume that $c(vw_3)=6$ and $c(vw_4)=7$. By Claim 11, there exists $a,b\in\{6,7\}$ such that $a\in C(w_1)$ and $b\in C(w_2)$.

By Claim 9 and Claim 10, for any $x\in C\backslash(C(u)\cup C(v))$, $\text{mult}_{S_v}(x)\geq 2$ and $\|S_v\cap C(u)\|\geq d(u)-2$. And recall that $a\in C(w_1)\cap\{6,7\}$, $b\in C(w_2)\cap\{6,7\}$, we have $\|S_v\|\geq 2|C\backslash(C(u)\cup C(v))|+|T_0|+(d(u)-2)+|\{a\}|+|\{b\}|=2(\Delta+11-(d(u)+d(v)))+|T_0|+(d(u)-2)+2=2\Delta+12+|T_0|-d(u)$. However, $\|S_v\|=\sum_{2\leq i\leq d(v)}d(v_i)-(d(v)-1)=\sum_{2\leq i\leq d(v)}d(v_i)-4\leq 2\Delta+d(v_2)+d(v_3)-4$. Clearly, $2\Delta+12+|T_0|-d(u)\leq 2\Delta+d(v_2)+d(v_3)-4$, which implies that $d(u)+d(v_2)+d(v_3)\geq 16+|T_0|$. Hence, $|T_0|\leq 3$ and $|T_0|=3$ if and only if $d(u)+d(v_2)+d(v_3)=19$ and $|C(u)\cap S_v|=d(u)-2$. Clearly, there exists $j_0\in\{\alpha_1,\alpha_2,\beta_1,\beta_2\}$ such that $\text{mult}_{S_v}(j_0)=2$ and assume that $j_0=\alpha_1\in T_2'$. Since $T_2\subseteq C(w_1)$, we can assume that $\alpha_1\in C(w_3)\backslash C(w_4)$. If H contains no $(6,\alpha_1)_{(w_2,w_3)}$-path, then recolor vw_2 with α_1 and reduce the proof to the Case 1. By $(*_{4.1})$ and Lemma 3, no bichromatic cycles are produced. Otherwise, we assume that:
$(*_{4.2})$ $6\in C(w_2)$ and H contains a $(6,\alpha_1)_{(w_2,w_3)}$-path, which cannot pass through w_4.

Claim 12

(a) *If* $|T_2'|\geq 2$, *then* $T_2'\subseteq C(w_3)\backslash C(w_4)$. *And for any* $\alpha_i\in T_2'$, H *contains a* $(6,\alpha_i)_{(w_2,w_3)}$-*path, which cannot pass through* w_4.

(b) *If* $|T_1'|\geq 1$, *then there exists* $j\in\{3,4\}$ *and* $k\in\{3,4\}\backslash\{j\}$ *such that* $T_1'\subseteq C(w_j)\backslash C(w_k)$. *And for any* $\beta_i\in T_1'$, H *contains a* $(c(vw_j),\beta_i)_{(w_1,w_j)}$-*path, which cannot pass through* w_k, *and* $c(vw_j)\in C(w_1)$.

Proof. To prove (a), we assume that there exists $\alpha_2 \in T_2'\backslash\{\alpha_1\}$ such that $\alpha_2 \in C(w_4)\backslash C(w_3)$. Then, we recolor vw_4 with α_1, vw_2 with α_2. Since $\alpha_1 \notin C(w_2)$, $\alpha_2 \notin C(w_3)$ and by $(*_{4.2})$ and Lemma 3, no bichromatic cycles are produced and reduce the proof to the Case 1. Thus, $T_2' \subseteq C(w_3)\backslash C(w_4)$. If H contains no $(6, \alpha_i)_{(w_2,w_3)}$-path for some $\alpha_i \in T_2'$, then recolor vw_2 with α_i and reduce the proof to the Case 1. Similarly, we can obtain (b). $\square$

By Claim 12, $T_2' \subseteq C(w_3)$. Since $\alpha_i \in C(w_3) \cap C(w_4)$ if $\mathrm{mult}_{S_v}(\alpha_i) \geq 3$ for any $\alpha_i \in T_2\backslash T_2'$, we have $T_2 \subseteq C(w_3)$. Thus, $d(w_3) \geq |T_2| + |\{6\}| = |T_2| + 1$. Similarly, there exists $j \in \{3, 4\}$ such that $T_1 \subseteq C(w_j)$ and $d(w_j) \geq |T_1| + |\{c(vw_j)\}| = |T_1| + 1$.

Claim 13 $C(u) \subseteq S_v$.

Proof. If $2 \notin S_v$, we recolor vw_3 with 2, vw_2 with α_1 and color uv with β_1. By $(*_{4.1})$ and Lemma 3, no bichromatic cycles are produced. Assume that $(C(u)\backslash\{1, 2\})\backslash S_v \neq \emptyset$ and $3 \in C(u)\backslash S_v$. If H contains no $(3, \beta_1)_{(u_3,w_2)}$-path, then recolor vw_2 with 3 and color uv with β_1. Otherwise, recolor vw_3 with 3, vw_2 with α_1 and color uv with β_1.

Assume that $1 \notin S_v$. If $|T_1'| \geq 1$, $\beta_1 \in T_1'$ and assume that $\beta_1 \in C(w_3)\backslash C(w_4)$ (it is similar if $\beta_1 \in C(w_4)\backslash C(w_3)$). Then, we recolor vw_4, vw_3, vw_1 with β_1, 1 and β_2, and color uv with α_1. Clearly, no bichromatic cycles are produced by $(*_{4.1})$ and Lemma 3. Otherwise, $|T_1'| = 0$ and $\mathrm{mult}_{S_v}(\beta_i) \geq 3$ for any $\beta_i \in T_1$, which implies that $|T_0| \geq |T_1|$. If H contains no $(6, \beta_1)_{(w_1,w_3)}$-path, then recolor vw_4 with 1, vw_1 with β_1 and color uv with α_1. Otherwise, H contains a $(6, \beta_1)_{(w_1,w_3)}$-path and $6 \in C(w_1)$. If H contains no $(7, \beta_1)_{(w_1,w_4)}$-path, then recolor vw_3 with 1, vw_1 with β_1 and color uv with α_1. Otherwise, H contains a $(7, \beta_1)_{(w_1,w_4)}$-path and $7 \in C(w_1)$. It follows that $1, 6, 7 \in C(w_1)$ and $|T_1| \geq \Delta + 11 - (d(u) + d(v)) - (d(w_1) - 3) = \Delta + 11 - (d(u) + d(v)) - d(w_1) + 3 \geq \Delta - d(w_1) + 3 \geq 3$ since $d(u) + d(v) \leq 11$ and $d(w_1) \leq \Delta$. Recall that $|T_1| \leq |T_0| \leq 3$. We have $|T_0| = |T_1| = 3$, $d(u) + d(v) = 11$, $d(w_1) = \Delta$ and $d(u) + d(v_2) + d(v_3) = 19$, which implies that $d(u) = d(v_2) = 6$, $d(v_i) = 7$ for $i = 3, 4, 5$ and $\Delta = 7$. And recall that $6, 7 \in C(w_1)$, $6 \in C(w_2)$, we have $\|S_v\| \geq 2|C\backslash(C(u) \cup C(v))| + |T_0| + (|C(u)| - 1) + \mathrm{mult}_{S_v}(6) + \mathrm{mult}_{S_v}(7)$ $= 2(\Delta + 11 - (d(u) + d(v))) + |T_0| + (d(u) - 2) + 3 = 2\Delta + 3 + 4 + 3 = 2 \times 7 + 10 = 24$. However, $\|S_v\| = \sum_{2 \leq i \leq d(v)} d(v_i) - (d(v) - 1) = 6 + 3 \times 7 - 4 = 23$, a contradiction. $\square$

Claim 14 *G contains a $((1, 7)_{(u_1,w_1)})$-path or a $(2, 7)_{(u_2,w_2)}$-path;* $\mathrm{mult}_{S_v}(6) + \mathrm{mult}_{S_v}(7) \geq 3$, *$7 \in C(w_2)$ or $7 \in C(w_1) \cap C(w_3)$ and H contains a $(6, 7)_{(w_2,w_3)}$-path.*

Proof. If H contains no $(1, 7)_{(u_1,w_1)}$-path and $(2, 7)_{(u_2,w_2)}$-path, then recolor vw_4 with α_1 and color uv with 7. By $(*_{4.2})$ and Lemma 3, no bichromatic cycles are produced. Otherwise, H contains a $(1, 7)_{(u_1,w_1)}$-path or $(2, 7)_{(u_2,w_2)}$-path and $7 \in C(w_1) \cup C(w_2)$. If $7 \in C(w_2)$, then $\mathrm{mult}_{S_v}(6) + \mathrm{mult}_{S_v}(7) \geq \|\{6, 7, a\}\| = 3$. Otherwise, $7 \in C(w_1)\backslash C(w_2)$ and H contains a $(1, 7)_{(u_1,w_1)}$-path. If H contains no $(6, 7)_{(w_2,w_3)}$-path, then recolor vw_2 with 7, vw_4 with α_1 and color uv with β_1. And no bichromatic cycles are produced by $(*_{4.2})$ and Lemma 3. Otherwise, H contains a $(6, 7)_{(w_2,w_3)}$-path and $7 \in C(w_3)$. Hence, $\mathrm{mult}_{S_v}(6) + \mathrm{mult}_{S_v}(7) \geq \mathrm{mult}_{C(w_1)}(7) + \mathrm{mult}_{C(w_2)}(6) + \mathrm{mult}_{C(w_3)}(7) = 3$. $\square$

Recall that $C(u) \subseteq S_v$ by Claim 13. Hence, $\|S_v\| \geq 2|C\backslash(C(u) \cup C(v))| + |T_0| + |C(u)| +$ $\text{mult}_{S_v}(6) +$ $\text{mult}_{S_v}(7) = 2(\Delta + 11 - (d(u) + d(v))) + |T_0| + d(u) - 1 + \text{mult}_{S_v}(6) + \text{mult}_{S_v}(7) \geq$ $2\Delta + 12 - d(u) + |T_0| - 1 + 3 = 2\Delta + 14 + |T_0| - d(u)$. However, $\|S_v\| = \sum_{2 \leq i \leq d(v)} d(v_i) -$ $(d(v) - 1) = \sum_{2 \leq i \leq d(v)} d(v_i) - 4 \leq 2\Delta + d(v_2) + d(v_3) - 4$. Hence, $2\Delta + 14 + |T_0| - d(u) \leq$ $2\Delta + d(v_2) + d(v_3) - 4$, which implies that $d(u) + d(v_2) + d(v_3) \geq 18 + |T_0|$.

Since $d(u) + d(v_2) + d(v_3) \leq 19$, we have:

$(*_{4.3})$ (i) $d(u) + d(v_2) + d(v_3) = 18$, $|T_0| = 0$ and $\text{mult}_{S_v}(6) + \text{mult}_{S_v}(7) = 3$; or (ii) $d(u) + d(v_2) + d(v_3) = 19$, $|T_0| \leq 1$ and $|T_0| = 1$ if and only if $\text{mult}_{S_v}(6) + \text{mult}_{S_v}(7) = 3$.

Let $\kappa_1 = |C(w_1) \cap (C(u) \cup C(v))|$, $\kappa_2 = |C(w_2) \cap (C(u) \cup C(v))|$. We have $|T_1| = |(C\backslash(C(u) \cup C(v)))\backslash C(w_1)| = \Delta + 11 - (d(u) + d(v)) - (d(w_1) - \kappa_1) \geq \Delta - d(w_1) + \kappa_1 \geq \kappa_1$ and $|T_2| = |(C\backslash(C(u) \cup C(v)))\backslash C(w_2)| = \Delta + 11 - (d(u) + d(v)) - (d(w_2) - \kappa_2) \geq \Delta - d(w_2) + \kappa_2 \geq \kappa_2$.

Claim 15 *If* $\text{mult}_{S_v}(6) +$ $\text{mult}_{S_v}(7) \geq 4$, *then*

(i) $|T_0| = 0$, *and* $d(u) = d(v_2) = 6$, $d(v_i) = 7$, $i = 3, 4, 5$;

(ii) $\text{mult}_{S_v}(6) +$ $\text{mult}_{S_v}(7) = 4$, *and for any* $i \in C(u)$, $\text{mult}_{S_v}(i) = 1$;

(iii) $|C\backslash(C(u) \cup C(v))| = \Delta$ *and* $|T_1| = \Delta - d(w_1) + \kappa_1$, $|T_2| = \Delta - d(w_2) + \kappa_2$.

(iv) *If* $|T_1| = \kappa_1$, *then* $\Delta = d(w_1) = 7$; *If* $|T_2| = \kappa_2$, *then* $\Delta = d(w_2) = 7$.

Proof. Since $\|S_v\| \geq 2|C\backslash(C(u) \cup C(v))| + |T_0| + \sum_{x \in C(u)} \text{mult}S_v(x) +$ $\text{mult}S_v(6) +$ $\text{mult}S_v(7)$ $\geq 2|C\backslash(C(u) \cup C(v))| + |T_0| + |C(u)| + 4 = 2(\Delta + 11 - (d(u) + d(v))) + |T_0| + d(u) - 1 + 4 \geq$ $2\Delta + 12 + |T_0| - d(u) + 3 = 2\Delta + 15 + |T_0| - d(u)$ and $\|S_v\| = \sum_{2 \leq i \leq d(v)} d(v_i) - (d(v) - 1) =$ $\sum_{2 \leq i \leq d(v)} d(v_i) - 4 \leq 2\Delta + d(v_2) + d(v_3) - 4$, we have $2\Delta + 15 + |T_0| - d(u) \leq 2\Delta + d(v_2) +$ $d(v_3) - 4$, which implies that $d(u) + d(v_2) + d(v_3) \geq 19 + |T_0|$. Since $|T_0| \geq 0$, we have $d(u) + d(v_2) + d(v_3) = 19$, $|T_0| = 0$. It follows that (ii)-(iv) holds. □

By Claim 12, Claim 15, we can further assume that:

$(*_{4.4})$ If $\text{mult}_{S_v}(6) + \text{mult}_{S_v}(7) \geq 4$, then $T_2' = T_2 \subseteq C(w_3)$, $T_1' = T_1 \subseteq C(w_j)$ for some $j \in \{3, 4\}$, $c(vw_j) \in C(w_1)$ and for any $\beta_i \in T_1$, H contains a $(c(vw_j), \beta_i)_{(w_1, w_j)}$-path, which cannot path through w_k, $k \in \{3, 4\}\backslash\{j\}$.

Since $|T_0| \leq 1$ by $(*_{4.3})$ and $|T_1| \geq 2$, we have $|T_1'| \geq 1$ and assume that $\beta_1 \in T_1'$. Now we need to consider the following.

Case 2.2.1 Assume that $\beta_1 \in C(w_3)$.

By Claim 12, $6 \in C(w_1)$. By Claim 14, $7 \in C(w_2)$ or $7 \in C(w_1) \cap C(w_3)$ and H contains a $(6, 7)_{(w_2, w_3)}$-path.

- $7 \in C(w_1) \cap C(w_2)$.

Then $\text{mult}_{S_v}(6) + \text{mult}_{S_v}(7) \geq 4$. By Claim 15, $d(w_3) \leq 7$, $6, 7 \notin C(w_3) \cup C(w_4)$, and for any $i \in C(u)$, $\text{mult}_{S_v}(i) = 1$, $T_1' = T_1 \subseteq C(w_3), T_2' = T_2 \subseteq C(w_3)$. Since $|T_1| \geq \lambda_1 \geq 3$, $|T_2| \geq \lambda_2 \geq 3$, $d(w_3) \leq 7$ and $d(w_3) \geq |T_1| + |T_2| + |\{6\}| \geq 7$, we have $T_1 = \{\beta_1, \beta_2, \beta_3\}$, $T_2 = \{\alpha_1, \alpha_2, \alpha_3\}$, $C(w_3) = \{6, \alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3\}$. We recolor vw_3 with 2, vw_2 with α_1 and color uv with β_1. By $(*_{4.1})$ and Lemma 3, no bichromatic cycles are produced.

- $7 \in C(w_2)\backslash C(w_1)$.

If H contains no $(6,7)_{(w_1,w_3)}$-path, we recolor vw_4 with α_1, vw_1 with 7 and color uv with α_2. By $(*_{4.1})$, $(*_{4.2})$ and Lemma 3, no bichromatic cycles are produced. Otherwise, H contains a $(6,7)_{(w_1,w_3)}$-path, which implies that $7 \in C(w_3)$. Thus, $\mathrm{mult}_{S_v}(6)+\mathrm{mult}_{S_v}(7) \geq 4$.

By Claim 15, $d(w_3) \leq 7$, $6 \notin C(w_4)$, and for any $i \in C(u)$, $\mathrm{mult}_{S_v}(i) = 1$, $T_1' = T_1 \subseteq C(w_3), T_2' = T_2 \subseteq C(w_3)$. Since $|T_1| \geq \kappa_1 \geq 2$, $|T_2| \geq \kappa_2 \geq 3$, and $d(w_3) \geq |T_1| + |T_2| + |\{6,7\}| \geq 7$, we have $T_1 = \{8,9\}$, $T_2 = \{10,11,12\}$, $C(w_3) = \{6,7,8,9,10,11,12\}$. Further, we may assume that $d(w_1) = d(w_2) = d(w_3) = \Delta = 7$ and $d(w_4) = 6$ by Claim 15, and $C(w_1) = \{1,6,10,11,12,13,14\}$, $C(w_2) = \{2,6,7,8,9,13,14\}$, $C(w_4) = \{1,2,3,4,5,7\}$. If H contains no $(1,13)_{(w_1,w_4)}$-path, recolor vw_4 with 13, vw_3 with 2, vw_2 with 10 and color uv with 8. Otherwise, H contains a $(1,13)_{(w_1,w_4)}$-path and $(2,13)_{(w_2,u)}$-path by $(*_{4.1})$, which cannot pass through w_4. Recolor vw_4, vw_1, vw_3 with 13, 8, 1, respectively, and color uv with 10. By $(*_{4.1})$ and Lemma 3, no bichromatic cycles are produced.

- $7 \in C(w_1)$ and $7 \in C(w_3) \backslash C(w_2)$.

Note that $\mathrm{mult}_{S_v}(6)+\mathrm{mult}_{S_v}(7) \geq 4$. By Claim 15, $d(w_3) \leq 7$, $6 \notin C(w_4)$, and for any $i \in C(u)$, $\mathrm{mult}_{S_v}(i) = 1$, $T_1' = T_1 \subseteq C(w_3), T_2' = T_2 \subseteq C(w_3)$. Since $|T_1| \geq 3$, $|T_2| \geq 2$, and $d(w_3) \geq |T_1| + |T_2| + |\{6,7\}| \geq 7$, we have $T_1 = \{10,11,12\}$, $T_2 = \{8,9\}$, $C(w_3) = \{6,7,8,9,10,11,12\}$. Further, we may assume that $d(w_1) = d(w_2) = d(w_3) = \Delta = 7$ and $d(w_4) = 6$ by Claim 15, and $C(w_1) = \{1,6,7,8,9,13,14\}$, $C(w_2) = \{2,6,10,11,12,13,14\}$, $C(w_4) = \{1,2,3,4,5,7\}$. If H contains no $(1,13)_{(w_1,w_4)}$-path, recolor vw_4 with 13, vw_3 with 2, vw_2 with 8 and color uv with 10. Otherwise, H contains a $(1,13)_{(w_1,w_4)}$-path and $(2,13)_{(w_2,u)}$-path by $(*_{4.1})$, which cannot pass through w_4. Recolor vw_4, vw_1, vw_3 with 13, 10, 1, respectively, and color uv with 8. By $(*_{4.1})$ and Lemma 3, no bichromatic cycles are produced.

Case 2.2.2 Assume that $\beta_1 \in C(w_4)$.

By Claims 12 and 14, $7 \in C(w_1)$ and if $7 \notin C(w_2)$, then $7 \in C(w_3)$, and H contains a $(6,7)_{(w_2,w_3)}$-path. If $6 \notin C(w_1)$ and G contains no $(2,6)_{(u_2,w_2)}$-path, then recolor ww_3 with β_1 and color uv with 6. Otherwise, G contains a $(2,6)_{(u_2,w_2)}$-path if $6 \notin C(w_1)$. If $6 \notin C(w_1)$ and H contains no $(6,7)_{(w_1,w_4)}$-path, then recolor vw_1 with 6, vw_3 with β_1 and color uv with α_1. By $(*_{4.4})$ and Lemma 3, no bichromatic cycles are produced. Otherwise, we assume that:

$(*_{4.5})$ (i) $7 \in C(w_2)$, or $7 \in C(w_3)$ and H contains a $(6,7)_{(w_2,w_3)}$-path. (ii) $6 \in C(w_1)$, or $6 \in C(w_4)$ and H contains a $(6,7)_{(w_1,w_4)}$-path.

By $(*_{4.5})$, we have $7 \in C(w_2) \cup C(w_3)$, $6 \in C(w_1) \cup C(w_4)$ and $\mathrm{mult}_{S_v}(6)+\mathrm{mult}_{S_v}(7) \geq 4$. By Claim 15 and $(*_{4.3})$, $(*_{4.4})$, $\mathrm{mult}_{S_v}(6)+\mathrm{mult}_{S_v}(7) = 4$, $d(w_i) \leq 7$ for any $1 \leq i \leq 4$, and for any $j \in C(u)$, $\mathrm{mult}_{S_v}(j) = 1$. By the previous analysis, we need to consider the following subcase.

- $c(vv_5) = 1$ and $7 \in C(v_5)$.

If $c(uv_5) = 2$, then we color uv with a color in $\{8, 9, \ldots, \Delta + 7\} \backslash C(v_5)$. Otherwise, $c(uv_5) = 3$. Assume that $c(vw_2) = 2$ and $c(vw_4) = 7$.

If there is no $(1,3)_{(v,w_2)}$-path in H, recolor vw_2 with 3 and color uv with a color in $\{8,9,\ldots,\Delta+7\}\backslash C(v_5)$. Since $\mathrm{mult}_{S_v}(3)=1$ and $3\in C(v_5)$, there is no bichromatic cycles. Otherwise, H contains a $(1,3)_{(v,w_2)}$-path and $1\in C(w_2)$. If for some color $i\in\{8,9,\ldots,\Delta+7\}\backslash C(v_5)$, there is no an $(i,j)_{(u,v_5)}$-path for any $j\in\{4,5\}$, then we recolor uv_5 with i and color uv with 3. Otherwise, for each color $i\in\{8,9,\ldots,\Delta+7\}\backslash C(v_5$, H contains an $(i,j)_{(u,v_5)}$-path for some color $j\in\{4,5\}$. Without loss of generality, assume that $4\in C(v_5)$ and H contains a $(\beta_1,4)_{(u,v_5)}$-path. Since $d(v_5)=7$ and $d(w_2)\le 7$, it suffice to assume that $C(v_5)=\{1,3,4,7,8,9,10\}$, $C(w_2)=\{1,2,6,11,12,13,14\}$ and $\Delta=7$. Then recolor vv_5, vw_2, vw_4 with $\beta_1,4,1$ and color uv with 8. By the previous assumption and Lemma 3, no bichromatic cycles are produced.

- $c(vv_5)=6$, $\beta_1\notin C(v_5)$ and $c(vw_1)=1$, $c(vw_2)=2$, $v(vw_4)=7$

If $c(uv_5)\in\{3,4,5\}$, then we recolor vw_2 with $c(uv_5)$ and color uv with β_1. Since $\mathrm{mult}_{S_v}(3)=1$, $3\notin C(v_5)$ and no bichormtic cycles are produced. Otherwise, $c(uv_5)\in\{1,2\}$. If for some color $i\in\{8,9,\ldots,\Delta+7\}\backslash C(v_5)$, there is no an $(i,j)_{(u,v_5)}$-path for any $j\in\{3,4,5\}$, then we recolor uv_5 with i and reduce the proof to the Case 1. Otherwise, for each color $i\in\{8,9,\ldots,\Delta+7\}\backslash C(v_5)$, H contains $(i,j)_{(u,v_5)}$-path for some color $j\in\{3,4,5\}$. Without loss of generality, assume that $3\in C(v_5)$ and H contains a $(\beta_1,3)_{(u,v_5)}$-path.

Assume that $c(uv_5)=1$. Since $\mathrm{mult}_{S_v}(3)=\mathrm{mult}_{S_v}(1)=1$, $1,3\notin(C(w_1)\backslash\{1\})\cup C(w_2)$. If H contains no $(3,6)_{(v_5,w_1)}$-path, then we recolor vw_1 with 3, vw_2 with 1 and color uv with β_1. Otherwise, H contains a $(3,6)_{(v_5,w_1)}$-path. Recolor vw_2 with 3, vv_5 with β_1 and color uv with 6. Since H contains no $(7,\beta_1)_{(w_1,w_4)}$-path by Claim 12, no bichromatic cycles are produced even $7\in C(v_5)$ by Lemma 3. Otherwise, assume that $c(uv_5)=2$. Since $\mathrm{mult}_{S_v}(3)=\mathrm{mult}_{S_v}(2)=1$, $2,3\notin C(w_1)$. If H contains no $(3,6)_{(v_5,w_1)}$-path, then we recolor vw_1 with 3 and color uv with β_1. Otherwise, H contains a $(3,6)_{(v_5,w_1)}$-path and $6\in C(w_1)$. Recolor vw_1 with 3, vv_5 with β_1 and color uv with 6. Since H contains no $(7,\beta_1)_{(w_1,w_4)}$-path by Claim 12, no bichromatic cycles are produced even $7\in C(v_5)$ by Lemma 3.

Case 3 $|C(u)\cap C(v)|=3$ and $C(u)\cap C(v)=\{1,2,3\}$.

Assume that $c(vw_3)=3$. By $(*_{4.1})$, for any $j\in C\backslash(C(u)\cup C(v))$, H contains an $(i,j)_{(u,v)}$-path for some $i\in\{1,2,3\}$ and $T_1\subseteq C_2\cup C_3\subseteq C(w_2)\cup C(w_3)$, $T_2\subseteq C_1\cup C_3\subseteq C(w_1)\cup C(w_3)$, and $T_3\subseteq C_1\cup C_2\subseteq C(w_1)\cup C(w_2)$. It is easy to derive that $|C\backslash(C(u)\cup C(v))|=\Delta+7-|C(u)\cup C(v)|=\Delta+7+2+3-(d(u)+d(v))=\Delta+12-(d(u)+d(v))\ge\Delta+1$, $|T_i|=|(C\backslash(C(u)\cup C(v)))\backslash C(w_i)|=|C\backslash(C(u)\cup C(v))|-(d(w_i)-1)\ge 2$, $i=1,2,3$.

Case 3.1 $(A_{4.1})$ holds and $d(v)=4$.

Assume that $1\notin S_v$. If H contains no $(2,\beta_1)_{(w_1,w_2)}$-path, then we recolor vw_3 with 1, vw_1 with β_1 and reduce the proof to the Case 2. Otherwise, H contains a $(2,\beta_1)_{(w_1,w_2)}$-path and a $(3,\beta_1)_{(u,w_3)}$-path by $(*_{4.1})$ and Lemma 3. Then we recolor vw_1 with β_1, vw_2 with 1 and reduce the proof to the Case 2. Otherwise, $1\in S_v$ and $2,3\in S_v$. Further, if $C(u)\backslash S_v\neq\emptyset$ and $4\in C(u)\backslash S_v$, we recolor vw_1 with 4 and then $4\in S_v$ similarly. Now we assume that $C(u)\subseteq S_v$. By Claim 9, for any $x\in C\backslash(C(u)\cup C(v))$, $\mathrm{mult}_{S_v}(x)\ge 2$ and $\|S_v\|\ge 2|C\backslash(C(u)\cup C(v))|$. So

$|S_v| \geq 2|C\backslash(C(u)\cup C(v))|+|C(u)| = 2(\Delta+7-(d(u)-1+3-3))+(d(u)-1) = 2\Delta+15-d(u)$. $\|S_v\| = \sum_{v_i\neq u\in N(v)} d(v_i) - (d(v)-1) = \sum_{v_i\neq u\in N(v)} d(v_i) - 3 \leq 2\Delta + d(v_2) - 3$. Hence, $2\Delta + 15 - d(u) \leq 2\Delta + d(v_2) - 3$, which implies that $d(u) + d(v_2) \geq 18$, a contradiction.

Case 3.2 $(A_{4.2})$ holds and $d(v) = 5$.

Assume that $c(w_4) = 6$. Recall that $|T_i| \geq 2$ for any $i \in \{1,2,3\}$.

Claim 16 *For* $i \in \{1,2,3\}$, $C(w_i)\cap C(v) \neq \{i\}$.

Proof. If $C(w_1)\cap C(v) = \{1\}$, then we recolor vw_1 with β_1 and reduce the proof to the Case 2. Hence, $C(w_1)\cap C(v) \neq \{1\}$ and assume that $\{1,a\} \subseteq C(v)\cap C(w_1)$. Similarly, we assume that $\{2,b\} \subseteq C(v)\cap C(w_2)$, $\{3,c\} \subseteq C(v)\cap C(w_3)$, where $a,b,c \in C(v)$. □

Claim 17 *For* $i = 1,2,3$, $T_i \subseteq C(w_j)$ *for any* $j \in \{1,2,3\}\backslash\{i\}$ *and* $T_i\cap T_j = \emptyset$ *for any* $i,j \in \{1,2,3\}$.

Proof. For any $\beta_i \in T_1\backslash T_1'$, it is clear that $\beta_i \in C(w_2)\cap C(w_3)$. If $T_1' \neq \emptyset$ and assume that $\beta_1 \in T_1'\backslash C(w_3)$, then H contains a $(2,\beta_1)_{(u,w_2)}$-path by $(*_{4.1})$ and $\beta_1 \in C(w_4)$ since $\mathrm{mult}_{S_v}(\beta_1) \geq 2$ by Claim 9. If H contains no $(\beta_1,6)_{(w_1,w_4)}$-path, we recolor vw_1 with β_1. Otherwise, H contains a $(\beta_1,6)_{(w_1,w_4)}$-path, recolor vw_3 with β_1. Clearly, no bichromatic cycles are produced by Lemma 3 and reduce the proof to the Case 2. Hence, $T_1' \subseteq T_1 \subseteq C(w_2)\cap C(w_3)$ and it is similar for $i = 2,3$. □

By Claims 16 and 17, $|T_i| \geq 3$ and $d(w_1) \geq |\{1,a\}|+|T_2|+|T_3|$, $d(w_2) \geq |\{2,b\}|+|T_1|+|T_3|$, $d(w_3) \geq |\{3,c\}|+|T_1|+|T_2|$.

Let $d(u) = 6-k$, $0 \leq k \leq 2$. Then $|C\backslash(C(u)\cup C(v))| = \Delta+7-(d(u)-1+d(v)-1)+|C(u)\cap C(v)| = \Delta+1+k$, $|T_i| \geq \Delta+1+k-(\Delta-2) \geq k+3$ and $d(w_i) \geq 2+2(k+3) = 2k+8$ for $i \in \{1,2,3\}$. Noting that $d(v_2) \geq d(u)$, we have $d(v_2)+d(v_3) \geq 2k+8+\min\{6-k, 2k+8\} = 2k+8+6-k = k+14$ and $d(u)+d(v_2)+d(v_3) \geq 6-k+k+14 = 20$, a contradiction. □

Case 4 $|C(u)\cap C(v)| = 4$ and $C(u)\cap C(v) = \{1,2,3,4\}$.

Then $d(v) = 5$ and assume that $c(vw_3) = 3$, $c(vw_4) = 4$. Clearly, $|C\backslash(C(u)\cup C(v))| \geq \Delta+2$ and for any $i \in \{1,2,3,4\}$, $|T_i| \geq 3$.

If $C(w_i)\cap C(v) = \{i\}$ for some $i \in \{1,2,3,4\}$, we recolor vw_i with a color in T_i and reduce the proof to the Case 3. Otherwise, for any $i \in \{1,2,3,4\}$, $C(w_i)\cap C(v) \neq \{i\}$. Assume that $\{1,a\} \subseteq C(w_1)\cap C(v)$, $\{2,b\} \subseteq C(w_2)\cap C(v)$, $\{3,c\} \subseteq C(w_3)\cap C(v)$, $\{4,d\} \subseteq C(w_4)\cap C(v)$.

If $d(u) = 5$, then $|C\backslash(C(u)\cup C(v))| \geq \Delta+3$ and $d(v_2)+d(v_3) \leq 18-5 = 13$. Hence, $2\Delta+6+4 \leq 2|C\backslash(C(u)\cup C(v))|+|\{a,b,c,d\}| \leq \|S_v\| = \sum_{2\leq i\leq d(v)} d(v_i)-(d(v)-1) = \sum_{2\leq i\leq d(v)} d(v_i)-4 \leq 2\Delta+d(v_2)+d(v_3)-4 \leq 2\Delta+13-4 = 2\Delta+9$, a contradiction.

If $d(u) = 6$, then $|C\backslash(C(u)\cup C(v))| \geq \Delta+2$ and $d(v_2)+d(v_3) \leq 19-6 = 13$. Hence, $2\Delta+4+4+|T_0| \leq 2|C\backslash(C(u)\cup C(v))|+|\{a,b,c,d\}|+|T_0| \leq \|S_v\| = \sum_{2\leq i\leq d(v)} d(v_i) - (d(v)-1) = \sum_{2\leq i\leq d(v)} d(v_i)-4 \leq 2\Delta+d(v_2)+d(v_3)-4 \leq 2\Delta+13-4 = 2\Delta+9$, i.e., $2\Delta+8+|T_0| \leq 2\Delta+9$, which implies that $|T_0| \leq 1$ and $|T_1'| \geq 1$ since $|T_1| \geq 4$. Assume that $\beta_1 \in T_1'$ and $\beta_1 \in (C(w_2) \cap C(w_3)) \backslash C(w_4)$. By $(*_{4.1})$, H contains a $(i, \beta_1)_{(u,v)}$-path for some $i \in \{2, 3\}$. Without loss of generality, assume that H contains a $(2, \beta_1)_{(u,v)}$-path. If H contains no $(3, \beta_1)_{(w_1,w_3)}$-path, we recolor vw_1 with β_1 and reduce the proof to the Case 3. Otherwise, recolor vw_4 with β_1 and reduce the proof to the Case 3. □

References

[1] N. Alon, B. Sudakov, and A. Zaks, *Acyclic edge colorings of graphs,* J. Graph Theory, 37 (2001), pp. 157-167.

[2] N. Alon, C. McDiarmid, and B. Reed, *Acyclic coloring of graphs*, Random Structures Algorithms, 2 (1991), pp. 277-288.

[3] M. Borowiecki and A. Fiedorowicz, *Acyclic edge coloring of planar graphs without short cycles*, Disc. Math., 310 (2010), pp. 1445-1455.

[4] M. Basavaraju and L. S. Chandran, *Acyclic edge coloring of subcubic graphs*, Disc. Math., 308 (2008), pp. 6650-6653.

[5] M. Basavaraju, L. S. Chandran, N. Cohen, F. Havet, and T. Müller, *Acyclic edge-coloring of planar graphs,* SIAM J. Disc. Math., 25 (2011), pp. 463-478.

[6] A. Fiedorowicz, M. Haluszczak, and N. Narayanan, *About acyclic edge colourings of planar graphs*, Inform. Process. Lett., 108 (2008), pp. 412-417.

[7] W. Dong and B. Xu, *Some results on acyclic edge coloring of plane graphs*, Inform. Process. Lett., 110 (2010), pp. 887-892.

[8] J. Fiamčik, *The acyclic chromatic class of a graph*, Math. Slovaca, 28 (1978), pp. 139-145. (in Russian)

[9] J. Hou, G. Liu, and J. Wu, *Acyclic edge coloring of planar graphs without small cycles*, Graphs Combin., DOI 10.1007/s00373-011-1043-0.

[10] J. Hou, J. Wu, G. Liu, and B. Liu, *Acyclic edge colorings of planar graphs and series-parallel graphs,* Sci. China Ser. A, 52 (2009), pp. 605-616.

[11] J. Hou, J. Wu, G. Liu, and B. Liu, *Acyclic edge chromatic number of outerplanar graphs*, J. Graph Theory, 64 (2010), pp. 22-36.

[12] M. Basavaraju and L. S. Chandran, *Acyclic edge coloring of graphs with maximum with degree 4*, J. Graph Theory, 61 (2009), pp. 192-209.

[13] R. Muthu, N. Narayanan, and C. R. Subramanian, *Acyclic edge colouring of outerplanar graphs*, Lecture Notes Comput. Sci., 4508 (2007), pp. 144-152.

[14] M. Molloy and B. Reed, *Further algorithmic aspects of Lovász local lemma*, in Proceedings of the 30th Annual ACM Symposium on Theory of Computing, 1998, pp. 524-529.

[15] J. Něsetřil and N. C. Wormald, *The acyclic edge chromatic number of a random d-regular graph is $d+1$*, J. Graph Theory, 49 (2005), pp. 69-74.

[16] S. Skulrattanakulchai, *Acyclic colorings of subcubic graphs*, Inform. Process. Lett., 92 (2004), pp. 161-167.

[17] Q. Shu and W. Wang, *Acyclic chromatic indices of planar graphs with girth at least five,* J. Comb. Optim., 23 (2012), pp. 140-157.

[18] W. Wang, Q. Shu, K. Wang, and P. Wang, *Acyclic chromatic indices of planar graphs with large girth,* Disc. Appl. Math., 59 (2011), pp. 1239-1253.

[19] W. Wang and K.-W. Lih, *Choosability and edge-choosability of planar graphs without* 5-*cycles,* Appl. Math. Lett., 15 (2002), pp. 561-565.

[20] Q. Shu, W. Wang, and Y. Wang, *Acyclic edge coloring of planar graphs without* 5-*cycles,* Disc. Appl. Math., DOI:10.1016/j.dam.2011.12.016.

[21] W. Wang and Q. Shu, *Acyclic chromatic indices of* K_4*-minor free graphs*, Sci. China Ser. A. 41 (2011), pp. 733-744.

[22] D. Yu, J. Hou, G. Liu, B. Liu, and L. Xu, *Acyclic Acyclic edge coloring of planar graphs with large girth*, Theoret. Comput. Sci., 410 (2009), pp. 5196-5200.

[23] V. Vizing, *On an estimate of the chromatic index of a* p*-graph*, Diskret Analiz. 3 (1964), pp. 25-30.